

Degenerate Parabolic Equation with Zero-flux Boundary Condition and its Approximations

MOHAMED KARIMOU GAZIBO

École Normale Supérieure, Département de Mathématiques
 Université Abdou Moumouni

Laboratoire de Mathématiques Fondamentales et Applications (LMFA)

BP:10963 Niamey
 NIGERIA

Abstract: We study a degenerate parabolic-hyperbolic equation with zero-flux boundary condition. The aim of this paper is to prove convergence of numerical approximate solutions towards the unique entropy solution. We propose an implicit finite volume scheme on admissible mesh. We establish fundamental estimates and prove that the approximate solution converge towards an entropy-process solution. Contrarily to the case of Dirichlet condition, in zero-flux problem unnatural boundary regularity of the flux is required to establish that entropy-process solution is the unique entropy solution. In the study of well-posedness of the problem, tools of nonlinear semigroup theory (stationary, mild and integral solutions) were used in order to overcome this difficulty. Indeed, in some situations including the one-dimensional setting, solutions of the stationary problem enjoy additional boundary regularity. Here, similar arguments are developed based on the new notion of integral-process solution that we introduce for this purpose.

Key-Words: Hyperbolic-parabolic equation, Finite volume scheme; Zero-flux boundary condition; Convergence, Boundary regularity, Entropy solution; Nonlinear semigroup theory; Mild solution; Integral-process solution.

Received: April 11, 2024. Revised: September 6, 2024. Accepted: September 26, 2024. Published: October 22, 2024.

1 Introduction

Let Ω be a bounded open set of \mathbb{R}^ℓ , $\ell \geq 1$, with a Lipschitz boundary $\partial\Omega$ and η the unit normal to $\partial\Omega$ outward to Ω . Let $Q = (0, T) \times \Omega$ and $\Sigma = (0, T) \times \partial\Omega$, T is a fixed positive time. We consider the zero-flux boundary problem

$$(P) \begin{cases} u_t + \operatorname{div} f(u) - \Delta\phi(u) = 0 & \text{in } Q \\ u(0, x) = u_0(x) & \text{in } \Omega, \\ (f(u) - \nabla\phi(u)) \cdot \eta(x) = 0 & \text{on } \Sigma. \end{cases}$$

The function f is continuous and satisfies:

$$f(0) = f(u_{\max}) = 0 \text{ for some } u_{\max} > 0. \quad (1)$$

We suppose that the initial data u_0 takes values in $[0, u_{\max}]$. In this case $[0, u_{\max}]$ will be an invariant domain for the solution of (P) in [1]. The function ϕ is non decreasing Lipschitz continuous in $[0, u_{\max}]$. Formally $\Delta(\phi(u)) = \operatorname{div}(\phi'(u)\nabla u)$. Then, if $\phi'(u) = 0$ for some $(t, x) \in Q$, the diffusion term vanishes so that (P) is a degenerate parabolic-hyperbolic problem. In our context, we suppose as in [1], that there exists a real value u_c with $0 \leq u_c \leq u_{\max}$ such that for $u \leq u_c$, the problem (P) is hyperbolic. This means that $\phi \equiv 0$ on $[0, u_c]$ and ϕ is strictly increasing in $[u_c, u_{\max}]$. Also as in [1], we assume that the couple (f, ϕ) is non-degenerate, this means that for all

$$\xi \in \mathbb{R}^\ell, \xi \neq 0, \text{ the functions } \lambda \mapsto \sum_{i=1}^{\ell} \xi_i f_i(\lambda)$$

are not affine on the non-degenerate sub intervals of $[0, u_c]$. It is well know that uniqueness of weak solution of degenerate hyperbolic-parabolic problem is not ensured, and we have to define a notion of entropy solution in the sense of [2], in the strictly hyperbolic case the fundamental paper of [3] to recover uniqueness. Inspired by [4], we defined in [1], a suitable notion of entropy solution for (P). A measurable function u taking values on $[0, u_{\max}]$ is called an entropy solution of the initial-boundary value problem (P) if $\phi(u) \in L^2(0, T; H^1(\Omega))$ and $\forall k \in [0, u_{\max}]$, $\forall \xi \in C^\infty([0, T) \times \mathbb{R}^\ell)^+$, the following inequality holds

$$\begin{aligned} & \int_0^T \int_{\Omega} |u - k| \xi_t dx dt + \int_{\Omega} |u_0 - k| \xi(0, x) dx \\ & + \int_0^T \int_{\Omega} \operatorname{sign}(u - k) \left[f(u) - f(k) - \nabla\phi(u) \right] \cdot \nabla \xi dx dt \\ & + \int_0^T \int_{\partial\Omega} |f(k) \cdot \eta(x)| \xi(t, x) d\mathcal{H}^{\ell-1}(x) dt \geq 0. \quad (2) \end{aligned}$$

In [5], the authors proved convergence of monotone scheme for a scalar conservation laws with zero-flux boundary condition which can be seen has a particular

case of our problem. Let us recall the main theoretical results on problem (P) obtained in [1]. We proved existence of solution satisfying (2), for any space dimension in the case $0 < u_c < u_{\max}$. Uniqueness is obtained for one space dimension. Remark that uniqueness is also true in multi-dimensional situation in two extreme cases: $u_c = 0$ for non-degenerate parabolic case [6], and $u_c = u_{\max}$ for pure hyperbolic case [4]. We refer to Appendix 2 for some explanations.

In this paper, we choose an implicit finite volume scheme for the discretization of the degenerate parabolic equation (P). Under suitable assumptions on the numerical fluxes, it is shown that the considered schemes are L^∞ stable and the discrete solutions satisfy some weak BV inequality and H^1 estimates. We prove also space and time translation estimates on the diffusion fluxes, which are the keys for the proof of convergence of the scheme. We prove existence of discrete solution by using Leray-Schauder topological degree. The approximate solutions are shown to satisfy an appropriate discrete entropy inequalities. Using the weak BV and H^1 estimates, the approximate solutions are also shown to satisfy continuous entropy inequalities. It remains to prove that the sequence of approximate solutions satisfying this continuous entropy inequalities converge towards a unique entropy solution. In [7], the authors use the concept of "entropy-process solution" introduced by [7], [8], [9], for Dirichlet boundary problem which is similar to the notion of measure valued solutions of [10]. They proved that approximate solutions converge towards an entropy-process solution as the mesh size tends to zero. Using doubling of variables method, they showed that the entropy-process solution is unique and is also a entropy solution of Dirichlet problem. In the case of zero-flux boundary condition, some difficulty due to lack of regularity of the boundary flux appears in [1]. We are not able to obtain uniqueness by the doubling of variables method. Thus, the only notion of entropy-process solution is not enough to prove convergence towards the entropy solution. To solve this difficulty, we found it useful to consider the general evolution problem of the form:

$$(E) \begin{cases} v'(t) + A(v(t)) = 0 & \text{on } (0, T) \\ v(0) = u_0. \end{cases}$$

We propose a new notion of solution called integral-process solution for the abstract evolution problem (E). This notion is presented in detail in the appendix 1. We prove that this new notion of integral-process solution coincides with the unique integral solution. Then, we apply this notion to the problem (P) and prove that the approximate solutions converge to an integral-process solution. We conclude that it is an entropy solution.

The rest of this paper is organized as follows. In Section 2, we present our implicit scheme. In Section 3, we prove a priori estimates, the discrete entropy inequalities and existence of discrete solution in Section 4. We propose in Section 5 a continuous entropy inequality, and the convergence result follows in Section 6. Finally, in Appendices we study the abstract evolution equation (E) and prove uniqueness of entropy solution in one space dimension for the degenerate parabolic problem (P).

2 Presentation of a finite volume scheme for degenerate parabolic problem with zero-flux boundary condition




Figure 1: Control volumes, centre, diamonds

In this section, we consider the problem (P) and construct a monotone finite volume scheme to approximate the solution. Let $\delta t > 0$ be the time step. Let \mathcal{O} be a family of disjoint connected polygonal subsets called control volumes of Ω such that $\bar{\Omega}$ is the union of the closures of the elements of this family and such that the common interface of two control volumes is included in the hyperplane of \mathbb{R}^ℓ . Let h be the upper bound for maximum size of the mesh: $h = \sup\{\text{Diam}(K), K \in \mathcal{O}\}$. We suppose that there exists $\alpha > 0$ such that:

$$\alpha h^\ell \leq m(K), m(\partial K) \leq \frac{1}{\alpha} h^{\ell-1}, \quad \forall K \in \mathcal{O}, \quad (3)$$

then the estimate on the number $|\mathcal{O}|$ of control volumes is

$$|\mathcal{O}| \leq \frac{m(\Omega)}{\alpha} h^{-\ell}, \quad (4)$$

where $m(K)$ is the ℓ -dimensional Lebesgue measure of K and $m(\partial K)$ is the $(\ell-1)$ -dimensional Lebesgue measure of ∂K . If K and L are two control volumes having an edge σ in common, we said that L is a neighbour of K and we write $L \in \mathcal{N}(K)$. We sometimes denote by $K|L$ the common edge σ between K and L and by $n_{K,\sigma}$ the unit normal to σ , oriented from K to L . Moreover, $\bar{\varepsilon}_K$ denotes the set of all edges for any control volumes K . If K has at least one common edge with boundary $\partial\Omega$, we denote by ε_K^{ext} the set of these boundaries edges, that can be regarded as $\varepsilon_K^{ext} = \{\sigma \in \bar{\varepsilon}_K, m(\sigma \cap \partial\Omega) > 0\}$. Eventually, if the control volume K has no common edges with a part of boundary $\partial\Omega$ then $\varepsilon_K^{ext} = \emptyset$. In all case, for all control volume $K \in \mathcal{O}$, we have $\varepsilon_K = \bar{\varepsilon}_K \setminus \varepsilon_K^{ext}$. Because we consider a zero-flux boundary condition, we don't need to distinguish between interior and exterior control volumes, only inner interfaces between volumes are needed in order to formulate the scheme. We consider here an admissible mesh of Ω for e.g., [7], [11], we mean that there exists a family of points $(x_K)_{K \in \mathcal{O}}$ such that the straight line $\overline{x_K x_L}$ is orthogonal to the interface $K|L$. We denote by $d_{K,L} = |x_K - x_L|$ the distance between x_K and x_L and by $d_{K,\sigma}$ the distance between x_K and the interface σ Figure 1. The point x_K is referred as the center of K . To simplify the analysis, we consider that $x_K \in K$ (in general, this assumption can be relaxed, e.g., one can consider so called Delaunay simplicial meshes). We denote by $\tau_{K,\sigma}$ the "transmissibility" through σ defined by $\tau_{K,\sigma} = \frac{m(\sigma)}{d_{K,\sigma}}$ if $\sigma \in \varepsilon_K^{ext}$, in addition we denote $\tau_{K|L} = \frac{m(K|L)}{d_{K,L}}$. The diamond denoted by $\widehat{K|L}$ is a convex hull constructed from neighbour centers x_K , x_L and $K|L$. The diamonds are disjoint and cover Ω up to an h -neighbourhood of $\partial\Omega$. Notice that the ℓ -dimensional measure $m(\widehat{K|L})$ of $\widehat{K|L}$ equals to $\frac{d_{K,L}}{\ell} m(K|L)$ Figure 1.

A discrete function w on the mesh \mathcal{O} is a set $(w_K)_{K \in \mathcal{O}}$. If w_K, v_K are discrete functions, the corresponding $L^2(\Omega)$ scalar product and norm can be computed as

$$(w_{\mathcal{O}}, v_{\mathcal{O}})_{L^2(\Omega)} = \sum_{K \in \mathcal{O}} m(K) w_K v_K;$$

$$\|w_{\mathcal{O}}\|_{L^2(\Omega)}^2 = \sum_{K \in \mathcal{O}} m(K) |w_K|^2.$$

In addition, we can define the positive (but not definite) product and the corresponding "discrete $H^1_{\mathcal{O}}$

semi-norm" by

$$\begin{aligned} & \ell \sum_{K \in \mathcal{O}} \sum_{L \in \mathcal{N}(K)} \frac{m(K|L)}{d_{K,L}} (w_L - w_K)(v_L - v_K) \\ & = (w_{\mathcal{O}}, v_{\mathcal{O}})_{H_{\mathcal{O}}} \end{aligned}$$

and

$$\|w_{\mathcal{O}}\|_{H_{\mathcal{O}}}^2 = (w_{\mathcal{O}}, w_{\mathcal{O}})_{H_{\mathcal{O}}}.$$

We define the discrete gradient $\nabla_{\mathcal{O}} w_{\mathcal{O}}$ of a constant per control volume function $w_{\mathcal{O}}$ as the constant per diamond $\widehat{K|L}$, \mathbb{R}^{ℓ} -valued function with values

$$(\nabla_{\mathcal{O}} w_{\mathcal{O}})_{\widehat{K|L}} = \nabla_{\widehat{K|L}} w_{\mathcal{O}} := \ell \frac{w_L - w_K}{d_{K,L}} \eta_{K,L}. \quad (5)$$

For the approximation of the convective term, we consider the numerical convection fluxes $F_{K,\sigma} : [0, u_{\max}]^2 \rightarrow \mathbb{R}$, $(a, b) \mapsto F_{K,\sigma}(a, b)$ for $K \in \mathcal{O}$, $\sigma \in \bar{\varepsilon}_K$

The numerical convection fluxes are monotone:

$$\partial_b F_{K,\sigma}(a, b) \leq 0 \leq \partial_a F_{K,\sigma}(a, b). \quad (6)$$

The numerical convection fluxes are conservative:

For all $\sigma = K|L$, for all $a, b \in [0, u_{\max}]$;

$$F_{K,L}(a, b) = -F_{L,K}(b, a). \quad (7)$$

The numerical convection fluxes are regular:

$F_{K,\sigma}$ is Lipschitz continuous and admits $m(\sigma)M$ as Lipschitz constant on $[0, u_{\max}]$.

$$(8)$$

The numerical convection fluxes are consistent:

$$\forall s \in [0, u_{\max}], F_{K,\sigma}(s, s) = m(\sigma) f(s) \cdot n_{K,\sigma}. \quad (9)$$

The Godunov, the splitting flux of Osher and Rusanov schemes may be the most common examples of schemes with fluxes satisfying (6)-(9).

Notice that the hypothesis (8) and (9) entail that for $\forall s \in [0, u_{\max}]$:

$$|F_{K,\sigma}(s, s)| \leq (\|f\|_{L^\infty} + u_{\max}) m(\sigma). \quad (10)$$

The discrete unknowns u_K^{n+1} for all control volume $K \in \mathcal{O}$, and $n \in \mathbb{N}$ are defined thanks to the following relations: first we initialize the scheme by

$$u_K^0 = \frac{1}{m(K)} \int_K u_0(x) dx \quad \forall K \in \mathcal{O}, \quad (11)$$

then, we use the implicit scheme for the discretization of problem (P):

$\forall n > 0, \forall K \in \mathcal{O}$,

$$m(K) \frac{u_K^{n+1} - u_K^n}{\delta t} + \sum_{\sigma \in \varepsilon_K} F_{K,\sigma}(u_K^{n+1}, u_{K,\sigma}^{n+1}) - \sum_{\sigma \in \varepsilon_K} \tau_{K,\sigma} \left(\phi(u_{K,\sigma}^{n+1}) - \phi(u_K^{n+1}) \right) = 0. \quad (12)$$

If the scheme has a solution, we will say that the piecewise constant function $u_{\mathcal{O},\delta t}(t, x)$ defined for $(t, x) \in]n\delta t, (n+1)\delta t] \times K$, a.e. by:

$$u_{\mathcal{O},\delta t}(t, x) = u_K^{n+1}. \quad (13)$$

is an approximate solution to (P).

Remark 2.1 1. Notice that using relation (9) and the fact that for all $s \in \mathbb{R} \operatorname{div}_x f(s) = 0$, we have $\forall s \in [0, u_{\max}], \forall K \in \mathcal{O}$:

$$\sum_{\sigma \in \varepsilon_K} F_{K,\sigma}(s, s) = 0. \quad (14)$$

This is equivalent to: $\forall s \in [0, u_{\max}], \forall K \in \mathcal{O}$

$$\sum_{\sigma \in \varepsilon_K} F_{K,\sigma}(s, s) + \sum_{\sigma \in \varepsilon_K^{ext}} F_{K,\sigma}(s, s) = 0. \quad (15)$$

2. Notice that the prescribed zero-flux boundary condition is in fact included in (12). One can extend the summation over $\sigma \in \bar{\varepsilon}_K$, and by convention regard the fluxes as:

$$\text{Flux}_{K,\sigma} = \begin{cases} \text{Flux}_{K,L} & \text{if } \sigma \in K|L, \\ 0 & \text{if } \sigma \in \varepsilon_K^{ext}. \end{cases}$$

3 Discrete entropy inequalities

This part is devoted to discrete entropy inequalities. We recall some notations in [8]:

Denote by $a \perp b = \min(a, b)$ and $a \top b = \max(a, b)$. We define $\eta_k^+(s) = (s-k)^+ = (s \top k - k)$, (respectively $\eta_k^-(s) = (s-k)^- = (s \perp k - k)$) and the associated fluxes-functions Φ_k^\pm called entropy fluxes

$$\begin{aligned} \Phi_k^+(s) &= \text{sign}^+(s-k)(f(s) - f(k)) \\ &= f(s \top k) - f(k); \\ \Phi_k^-(s) &= \text{sign}^-(s-k)(f(s) - f(k)) \\ &= f(s \perp k) - f(k); \\ \Phi_k(s) &= \text{sign}(s-k)(f(s) - f(k)). \end{aligned}$$

Therefore, the numerical sub and super entropy fluxes functions are defined by the formulas

$$\begin{aligned} \Phi_{K,\sigma,k}^+(a, b) &= F_{K,\sigma}(a \top k, b \top k) - F_{K,\sigma}(k, k); \\ \Phi_{K,\sigma,k}^-(a, b) &= F_{K,\sigma}(k, k) - F_{K,\sigma}(a \perp k, b \perp k); \\ \Phi_{K,\sigma,k}(a, b) &= F_{K,\sigma}(a \top k, b \top k) - F_{K,\sigma}(a \perp k, b \perp k). \end{aligned}$$

From now, we have the following the discrete entropy inequalities.

Lemma 3.1 Assume that (4), (6)- (9) hold. Let $u_{\mathcal{O},\delta t}$ be an approximate solution of the problem (P) defined by (11), (12). Then for all $k \in [0, u_{\max}]$, for all $K \in \mathcal{O}$, $n \geq 0$ the following discrete sub-entropy inequalities hold:

$$\begin{aligned} & \frac{\eta_k^+(u_K^{n+1}) - \eta_k^+(u_K^n)}{\delta t} m(K) \\ & + \sum_{\sigma \in \varepsilon_K} \Phi_{K,\sigma,k}^+(u_K^{n+1}, u_{K,\sigma}^{n+1}) \\ & - \sum_{K|L} \tau_{K|L} \left(\eta_{\phi(k)}^+(\phi(u_L^{n+1})) - \eta_{\phi(k)}^+(\phi(u_K^{n+1})) \right) \\ & \leq \sum_{\sigma \in \varepsilon_K^{ext}} \text{sign}^+(u_K^{n+1} - k) m(\sigma) f(k) n_{K,\sigma}. \quad (16) \end{aligned}$$

Also the discrete super-entropy inequalities are satisfied (i.e., η_k^+ , $\Phi_{K,\sigma,k}^+$, sign^+ can be replaced by η_k^- , and $\Phi_{K,\sigma,k}^-$, sign^- in (16)).

Notice that, if for all $K \in \mathcal{O}$, u_K^{n+1} satisfy both discrete sub-entropy inequality and discrete super-entropy inequality, then u_K^{n+1} can be seen as a discrete entropy solution in $]n\delta t, (n+1)\delta t] \times K$.

Proof. Thanks to the Remark 2.1, the constant $k \in [0, u_{\max}]$ is solution of:

$$\begin{aligned} & m(K) \frac{k - k}{\delta t} + \sum_{\sigma \in \varepsilon_K} F_{K,\sigma}(k, k) \\ & - \sum_{K|L} \tau_{K|L} \left(\phi(k) - \phi(k) \right) \\ & = - \sum_{\sigma \in \varepsilon_K^{ext}} F_{K,\sigma}(k, k). \quad (17) \end{aligned}$$

Subtracting from the equality (12) the equality (17), we obtain:

$$\begin{aligned} & \frac{1}{\delta t} \left((u_K^{n+1} - k) - (u_K^n - k) \right) m(K) \\ & + \sum_{\sigma \in \varepsilon_K} \left(F_{K,\sigma}(u_K^{n+1}, u_{K,\sigma}^{n+1}) - F_{K,\sigma}(k, k) \right) \\ & - \sum_{L \in N(K)} \tau_{K|L} \left[\left(\phi(u_L^{n+1}) - \phi(k) \right) - \left(\phi(u_K^{n+1}) - \phi(k) \right) \right] \\ & = \sum_{\sigma \in \varepsilon_K^{ext}} F_{K,\sigma}(k, k). \quad (18) \end{aligned}$$

Multiply (18) by $(\eta_k^+)'(u_K^{n+1}) = \text{sign}^+(u_K^{n+1} - k)$. We recall that for all convex function J , we have for all

$z_1, z_2 \in \mathbb{R}$, the convexity inequality $(z_1 - z_2)J'(z_1) \geq J(z_1) - J(z_2)$. (Here, we may consider J' as being multivalued, in the sense of sub-differential of J). First, we use this convexity inequality to obtain

$$\begin{aligned} & \text{sign}^+(u_K^{n+1} - k) \left((u_K^{n+1} - k) - (u_K^n - k) \right) \\ & \geq \left((u_K^{n+1} - k)^+ - (u_K^n - k)^+ \right). \end{aligned} \quad (19)$$

Second, due to the monotony of the numerical fluxes, we see that

$$\begin{aligned} & \text{sign}^+(u_K^{n+1} - k) \left(F_{K,\sigma}(u_K^{n+1}, u_{K,\sigma}^{n+}) - F_{K,\sigma}(k, k) \right) \\ & \geq \Phi_{K,\sigma,k}^+(u_K^{n+1}, u_{K,\sigma}^{n+}). \end{aligned} \quad (20)$$

Finally, using the convexity inequality and the monotonicity of ϕ , we have:

$$\begin{aligned} & -(\eta_k^+)'(u_K^{n+1}) \left(\phi(u_L^{n+1}) - \phi(u_K^{n+1}) \right) \\ & \geq -\left(\eta_{\phi(k)}^+(\phi(u_L^{n+1})) - \eta_{\phi(k)}^+(\phi(u_K^{n+1})) \right). \end{aligned} \quad (21)$$

Then, we get from (19), (20), (21)

$$\begin{aligned} & \frac{1}{\delta t} \left((u_K^{n+1} - k)^+ - (u_K^n - k)^+ \right) m(K) \\ & + \sum_{\sigma \in \mathcal{E}_K} \Phi_{K,\sigma,k}^+(u_K^{n+1}, u_{K,\sigma}^{n+}) \\ & - \sum_{K|L} \tau_{K|L} \left(\eta_{\phi(k)}^+(\phi(u_L^{n+1})) - \eta_{\phi(k)}^+(\phi(u_K^{n+1})) \right) \\ & \leq \sum_{\sigma \in \mathcal{E}_K^{ext}} \text{sign}^+(u_K^{n+1} - k) m(\sigma) f(k) n_{K,\sigma}. \end{aligned}$$

This prove (16). In the same way, we prove the discrete super-entropy inequalities. Finally, we deduce that u_K^{n+1} satisfies the discrete entropy inequality in this sense:

$$\begin{aligned} & \frac{\eta_k(u_K^{n+1}) - \eta_k(u_K^n)}{\delta t} m(K) + \sum_{\sigma \in \mathcal{E}_K} \Phi_{K,\sigma,k}(u_K^{n+1}, u_{K,\sigma}^{n+}) \\ & - \sum_{K|L} \tau_{K|L} \left(\eta_{\phi(k)}(\phi(u_L^{n+1})) - \eta_{\phi(k)}(\phi(u_K^{n+1})) \right) \\ & \leq \sum_{\sigma \in \mathcal{E}_K^{ext}} \text{sign}(u_K^{n+1} - k) m(\sigma) f(k) n_{K,\sigma}. \end{aligned} \quad (22)$$

4 Estimates of discrete solution and existence

We wish to prove that the approximate solution $u_{O,\delta t}$ satisfies the continuous entropy inequalities in section

5. To this purpose, we give fundamental estimates useful for proving convergence of the scheme. First, we prove the L^∞ stability of the scheme, this comes from discrete entropy inequalities and the L^∞ norm on the flux f with the relation (1).

4.1 L^∞ bound on discrete solutions

Proposition 4.1 *Suppose that $K \in \mathcal{O}$, the assumptions (4), (6)- (9) hold. Assume that $u_0 \in [0, u_{\max}]$. Then the approximate solution $u_{O,\delta t}(t, x)$ of problem (P) defined by (11), (12) satisfies:*

$$0 \leq u_K^n(t, x) \leq u_{\max} \quad \forall K \in \mathcal{O}. \quad (23)$$

Proof. Summing (16) over $K \in \mathcal{O}$, we get

$$\begin{aligned} & \sum_{K \in \mathcal{O}} m(K) \frac{\eta_k^+(u_K^{n+1}) - \eta_k^+(u_K^n)}{\delta t} \\ & + \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K} \Phi_{K,\sigma,k}^+(u_K^{n+1}, u_{K,\sigma}^{n+}) \\ & - \sum_{K \in \mathcal{O}} \sum_{L \in \mathcal{N}(K)} \tau_{K|L} \left(\eta_{\phi(k)}^+(\phi(u_L^{n+1})) - \eta_{\phi(k)}^+(\phi(u_K^{n+1})) \right) \\ & \leq \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K^{ext}} m(\sigma) |f(k) n_{K,\sigma}|. \end{aligned} \quad (24)$$

In inequality (24), take $k = u_{\max}$ and use (1) to obtain:

$$\begin{aligned} & \sum_{K \in \mathcal{O}} \frac{m(K)}{\delta t} \left(u_K^{n+1} - u_{\max} \right)^+ - \sum_{K \in \mathcal{O}} \frac{m(K)}{\delta t} \left(u_K^n - u_{\max} \right)^+ \\ & + \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K} \Phi_{K,\sigma,u_{\max}}^+(u_K^{n+1}, u_{K,\sigma}^{n+}) \\ & - \sum_{K \in \mathcal{O}} \sum_{L \in \mathcal{N}(K)} \tau_{K|L} \left(\eta_{\phi(u_{\max})}^+(\phi(u_L^{n+1})) - \eta_{\phi(u_{\max})}^+(\phi(u_K^{n+1})) \right) \\ & \leq 0. \end{aligned}$$

From now, remark that due to the conservativity of the scheme we have

$$\begin{aligned} & \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K} \Phi_{K,\sigma,u_{\max}}^+(u_K^{n+1}, u_{K,\sigma}^{n+}) \\ & = \sum_{K \in \mathcal{O}} \sum_{L \in \mathcal{N}(K)} \Phi_{K,L,u_{\max}}^+(u_K^{n+1}, u_L^{n+1}) = 0 \\ & \sum_{K \in \mathcal{O}} \sum_{L \in \mathcal{N}(K)} \left(\eta_{\phi(u_{\max})}^+(\phi(u_L^{n+1})) - \eta_{\phi(u_{\max})}^+(\phi(u_K^{n+1})) \right) = 0. \end{aligned}$$

Therefore

$$\sum_{K \in \mathcal{O}} \frac{m(K)}{\delta t} \left(u_K^{n+1} - u_{\max} \right)^+ - \left(u_K^n - u_{\max} \right)^+ \leq 0.$$

Since $0 \leq u_K^0 \leq u_{\max}$, by induction we prove $(u_K^{n+1} - u_{\max})^+ \leq 0$. In the same way, in the super-entropy inequality, taking $k = 0$, using (1), we also prove that $(u_K^{n+1})^- \leq 0$.

4.2 Weak BV and $L^2(0, T, H^1(\Omega))$ estimates

Now, we give the weak BV and $L^2(0, T, H^1(\Omega))$ estimates. The $L^2(0, T, H^1(\Omega))$ as the L^∞ estimate are necessary for justifying compactness properties of discrete solutions. The weak BV-stability does not give directly any compactness result, however, it plays a crucial role in the proof of continuous entropy inequality in section 5. To start with, we recall a Lemma which is one ingredient of the proof of Lemma 4.3 below.

Lemma 4.2 *Let $G : [a, b] \rightarrow \mathbb{R}$ be a monotone Lipschitz continuous function with Lipschitz constant $L > 0$ and $a, b \in \mathbb{R}$. Then for all $c, d \in [a, b]$, one has*

$$\left| \int_c^d (G(x) - G(c)) dx \right| \geq \frac{1}{2L} (G(d) - G(c))^2.$$

Proof. In order to prove this result, we assume, for instance, that G is nondecreasing and $c < d$ (the other cases are similar). Then, we have $G(s) \geq H(s)$, for all $s \in [c, d]$, where $H(s) = G(c)$ for $s \in [c, d - l]$ and $H(s) = G(c) + (s - d + l)L$ for $s \in [d - l, d]$, with $lL = G(d) - G(c)$, and therefore:

$$\begin{aligned} \int_c^d (G(s) - G(c)) ds &\geq \int_c^d (H(s) - G(c)) ds \\ &= \frac{l}{2} (G(d) - G(c)) \\ &= \frac{1}{2L} (G(d) - G(c))^2. \end{aligned}$$

Now, we establish the weak BV-stability of the scheme.

Lemma 4.3 (Weak BV-Estimate) *Suppose that (4),(6)-(9) hold. Let $u_{O, \delta t}$ be an approximate solution of problem (P) defined by (11), (12). Let $T > 0$, and set $N = \max\{n \in \mathbb{N}, n < \frac{T}{\delta t}\}$ and $L \in \mathcal{N}(K)$ (with convention $u_K^{n+1} \geq u_L^{n+1}$). Then there exists $C = C(\|f\|_{L^\infty}, u_{\max}, T, |\Omega|) \geq 0$ such that*

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_{K|L} \max_{u_L^{n+1} \leq c \leq d \leq u_K^{n+1}} (F_{K, \sigma}(d, c) - F_{K, \sigma}(d, d)) \\ &+ \sum_{n=0}^N \delta t \sum_{K|L} \max_{u_L^{n+1} \leq c \leq d \leq u_K^{n+1}} (F_{K, \sigma}(d, c) - F_{K, \sigma}(c, c)) \\ &\leq \frac{C}{\sqrt{h}}. \end{aligned} \tag{25}$$

Proof. Multiplying (12) by $\delta t u_K^{n+1}$ and summing over $K \in \mathcal{O}$ and $n = 0, \dots, N$ yields $A_{Ev} + A_{Cv} + A_{Dif} = 0$

with

$$\begin{aligned} A_{Ev} &= \sum_{n=0}^N \sum_{K \in \mathcal{O}} m(K) (u_K^{n+1} - u_K^n) u_K^{n+1}; \\ A_{Cv} &= \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K} F_{K, \sigma} (u_K^{n+1}, u_{K, \sigma}^{n+1}) u_K^{n+1}; \\ A_{Dif} &= - \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} \sum_{K|L} \tau_{K|L} (\phi(u_L^{n+1}) - \phi(u_K^{n+1})) u_K^{n+1} \end{aligned}$$

Let us first estimate A_{Ev} . We use the fact that:

$$\forall a, b \in \mathbb{R}, (a - b)a = \frac{1}{2}(a - b)^2 + \frac{1}{2}a^2 - \frac{1}{2}b^2,$$

we get:

$$\begin{aligned} A_{Ev} &= \sum_{n=0}^N \sum_{K \in \mathcal{O}} m(K) (u_K^{n+1} - u_K^n) u_K^{n+1} \\ &= \frac{1}{2} \sum_{n=1}^{N-1} \sum_{K \in \mathcal{O}} m(K) (u_K^{n+1} - u_K^n)^2 \\ &\quad + \frac{1}{2} \sum_{K \in \mathcal{O}} m(K) \left[(u_K^{N+1})^2 - (u_K^0)^2 \right]. \end{aligned} \tag{26}$$

The two first terms are non negative and due to (23) there exists $C \geq 0$ (that only depends on $|\Omega|$ and u_{\max}) such that $-C$ is a lower bound for the last term, then

$$A_{Ev} \geq -C. \tag{27}$$

Secondly, using summation by parts then A_{Dif} gives

$$\sum_{n=0}^N \delta t \sum_K \sum_{K|L} \tau_{K|L} (\phi(u_L^{n+1}) - \phi(u_K^{n+1})) (u_L^{n+1} - u_K^{n+1})$$

and $A_{Dif} \geq 0$. Now, we study the term A_{Cv} . Due to (15), it can be rewritten as the sum between A_{Cv}^{int} and A_{Cv}^{ext} :

$$\begin{aligned} &\sum_{n=0}^N \sum_{K|L} \delta t (F_{K, \sigma}(u_K^{n+1}, u_L^{n+1}) - F_{K, \sigma}(u_K^{n+1}, u_K^{n+1})) u_K^{n+1} \\ &- \sum_{n=0}^N \sum_{K|L} \delta t (F_{K, \sigma}(u_K^{n+1}, u_L^{n+1}) - F_{K, \sigma}(u_L^{n+1}, u_L^{n+1})) u_L^{n+1} \\ &= A_{Cv}^{int} \end{aligned}$$

$$\begin{aligned} A_{Cv}^{ext} &= - \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} u_K^{n+1} \sum_{\sigma \in \mathcal{E}_K^{ext}} F_{K, \sigma}(u_K^{n+1}, u_K^{n+1}) \\ &= - \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} u_K^{n+1} \sum_{\sigma \in \mathcal{E}_K^{ext}} m(\sigma) f(u_K^{n+1}) \cdot \eta_{K, \sigma}. \end{aligned} \tag{28}$$

We can estimate the boundary term A_{Cv}^{ext} by

$$|A_{Cv}^{ext}| \leq C(\|f\|_{L^\infty}, u_{\max}, T, |\partial\Omega|). \quad (29)$$

Let us assign:

$$\begin{aligned} \Psi_{K,L}(a) &= \int_0^a s \left(\frac{\partial F_{K,L}}{\partial u}(s, s) + \frac{\partial F_{K,L}}{\partial v}(s, s) \right) ds \\ &= \int_0^a s \frac{d}{ds} F_{K,L}(s, s) ds. \end{aligned}$$

Then

$$\begin{aligned} &\int_0^b s \frac{d}{ds} F_{K,L}(s, s) ds - \int_0^a s \frac{d}{ds} F_{K,L}(s, s) ds, \\ &= b \left(F_{K,L}(b, b) - F_{K,L}(a, b) \right) \\ &\quad - a \left(F_{K,L}(a, a) - F_{K,L}(a, b) \right) \\ &\quad - \int_a^b \left(F_{K,L}(s, s) - F_{K,L}(a, b) \right) ds \\ &= \Psi_{K,L}(b) - \Psi_{K,L}(a). \end{aligned} \quad (30)$$

Take $a = u_K^{n+1}$ and $b = u_L^{n+1}$ in (30) and multiply by δt . Summing over $n = 0, \dots, N$ and $L \in \mathcal{N}(K)$, we obtain $A_{Cv}^{int} = A_{Cv}^1 + A_{Cv}^2$, where:

$$\begin{aligned} A_{Cv}^1 &= \sum_{n=0}^{n+1} \sum_{K|L} \delta t \int_{u_L^{n+1}}^{u_K^{n+1}} \left(F_{K,L}(u_K^{n+1}, u_L^{n+1}) - F_{K,L}(s, s) \right) ds \\ A_{Cv}^2 &= - \sum_{n=0}^{n+1} \sum_{K|L} \delta t \left(\Psi_{K,L}(u_K^{n+1}) - \Psi_{K,L}(u_L^{n+1}) \right). \end{aligned}$$

We have $\sum_{\sigma \in \bar{\varepsilon}_K} \Psi_{K,L}(s) = 0$, for all $s \in [0, u_{\max}]$; then

it appears that A_{Cv}^2 reduces to the sum of $\sigma \in \varepsilon_K^{ext}$, and it satisfies $|A_{Cv}^2| \leq C$ similar to the estimate of A_{Cv}^{ext} .

Now, consider $a, b, c, d \in \mathbb{R}$ such that $a \leq c \leq d \leq b$. Take $i_\alpha := u_L^{n+1} \leq c \leq d \leq u_K^{n+1}$. Using the monotonicity of $F_{K,L}$ and Lemma 4.2, we deduce

$$\begin{aligned} &\int_a^b \left(F_{K,L}(b, a) - F_{K,L}(s, s) \right) ds \\ &\geq \int_d^c \left(F_{K,L}(d, c) - F_{K,L}(d, s) \right) ds \\ &\geq \frac{1}{2M} \left(F_{K,L}(d, c) - F_{K,L}(c, c) \right)^2. \end{aligned}$$

with $\bar{M} = m(K|L)M$. Take

$$I = \int_{u_L^{n+1}}^{u_K^{n+1}} \left(F_{K,L}(u_K^{n+1}, u_L^{n+1}) - F_{K,L}(s, s) \right) ds.$$

Therefore, we get:

$$\begin{aligned} I &\geq \frac{1}{2M} \max_{i_\alpha} \left(F_{K,L}(d, c) - F_{K,L}(d, d) \right)^2 \\ I &\geq \frac{1}{2M} \max_{i_\alpha} \left(F_{K,L}(d, c) - F_{K,L}(c, c) \right)^2. \end{aligned}$$

Then, we have

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_{K|L} \frac{1}{4M} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(d, d) \right)^2 \\ &\quad + \sum_{n=0}^N \delta t \sum_{K|L} \frac{1}{4M} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(c, c) \right)^2 \\ &\leq A_{Cv}^1. \end{aligned}$$

Recalling the equality $A_{Cv} = A_{Cv}^{ext} + A_{Cv}^{int,1} + A_{Cv}^{int,2}$, we find

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_{K|L} \frac{1}{4M} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(d, d) \right)^2 \\ &\quad + \sum_{n=0}^N \delta t \sum_{K|L} \frac{1}{4M} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(c, c) \right)^2 \\ &\quad - 2C \leq A_{Cv}. \end{aligned} \quad (31)$$

Set

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_{K|L} \frac{1}{4M} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(d, d) \right)^2 \\ &\quad + \sum_{n=0}^N \delta t \sum_{(K,L) \in \varepsilon_{int}^{n+1}} \frac{1}{4M} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(c, c) \right)^2 \\ &= \bar{A}_{Cv}. \end{aligned}$$

Now, as the equality $A_{Ev} + A_{Cv} + A_{Dif} = 0$ holds and as (27) and (32) are satisfied, we have $\bar{A}_{Cv} \leq C$.

Moreover, using the Cauchy-Schwarz inequality, we deduce

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_{K|L} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(d, d) \right) \\ &\quad + \sum_{n=0}^N \delta t \sum_{K|L} \max_{i_\alpha} \left(F_{K,\sigma}(d, c) - F_{K,\sigma}(c, c) \right) \\ &\leq \frac{1}{4M} \sqrt{\sum_{n=0}^N \delta t \sum_{K|L} m(K|L) \sqrt{\bar{A}_{Cv}}}. \end{aligned} \quad (32)$$

At the end, take into account the regularity on the mesh (4) to deduce that:

$$\begin{aligned} \sum_{n=0}^N \delta t \sum_{K|L} m(K|L) &\leq T \frac{1}{\alpha} h^{\ell-1} \sum_{K|L} 1 \\ &\leq T \frac{1}{\alpha} h^{\ell-1} m(\Omega) \frac{1}{\alpha} h^{-\ell} \leq \frac{C}{h}. \end{aligned}$$

Lemma 4.4 ($L^2(0, T, H^1(\Omega))$ Estimate) Suppose that (4), (6)-(9) hold. Let $u_{O, \delta t}$ be the approximate solution of problem (P) defined by (11), (12). Let $T > 0$, and set $N = \max\{n \in \mathbb{N}, n < \frac{T}{\delta t}\}$. Then there exists $C = C(\|f\|_{L^\infty}, u_{\max}, T) \geq 0$ such that

$$\frac{1}{2} \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in N(K)} \tau_{K|L} \left| \phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right|^2 \leq C. \quad (33)$$

Proof. Multiplying (12) by $\delta t \phi(u_K^{n+1})$ and summing over $K \in O$ and $n = 0, \dots, N$:

$$\begin{aligned} &\sum_{n=0}^N \sum_{K \in O} m(K) (u_K^{n+1} - u_K^n) \phi(u_K^{n+1}) \\ &- \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in N(K)} \tau_{K|L} \left(\phi(u_L^{n+1}) - \phi(u_K^{n+1}) \right) \phi(u_K^{n+1}) \\ &+ \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{\sigma \in \mathcal{E}_K} F_{K, \sigma} (u_K^{n+1}, u_{K, \sigma}^{n+1}) \phi(u_K^{n+1}) = 0. \end{aligned}$$

i.e $B_{Ev} + B_{Cv} + B_{Dif} = 0$.

Let $\vartheta(r) = \int_0^r \phi(s) ds$. From the convexity inequality, we have:

$$\begin{aligned} B_{Ev} &\geq \sum_{n=0}^N \sum_{K \in O} m(K) \left(\vartheta(u_K^{n+1}) - \vartheta(u_K^n) \right) \\ &= \sum_{K \in O} m(K) \left(\vartheta(u_K^{N+1}) - \vartheta(u_K^0) \right). \quad (34) \end{aligned}$$

Further, in the term B_{Dif} , for every edge $K|L$ the terms involving K and L appear twice. Thanks to the conservativity of the scheme, we find

$$B_{Dif} = - \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in N(K)} \tau_{K|L} \left| \phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right|^2. \quad (35)$$

The term B_{Cv} can be rewritten as

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_K \sum_L F_{K, L} (u_K^{n+1}, u_L^{n+1}) \left(\phi(u_L^{n+1}) - \phi(u_K^{n+1}) \right) \\ &= B_{Cv}. \end{aligned}$$

Using the weighted Young inequality and (10), we can major $|B_{Cv}|$ as follows

$$\begin{aligned} &\sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in N(K)} \frac{d_{K, L}}{2m(K|L)} \left(F_{K, \sigma} (u_K^{n+1}, u_L^{n+1}) \right)^2 \\ &+ \frac{1}{2} \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in N(K)} \frac{m(K|L)}{d_{K, L}} \left| \phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right|^2 \\ &\leq C \sum_{L|K} m(\widehat{K|L}) \\ &+ \frac{1}{2} \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in N(K)} \tau_{K|L} \left| \phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right|^2. \end{aligned}$$

Collecting the previous inequalities we readily deduce (33). This concludes the proof of the Lemma 4.4.

4.3 Estimates of space and time translates

Recall the following result.

Theorem 4.5 (Riesz-Frechet-Kolmogorov) Let an open $Q \subset \mathbb{R}^{\ell+1}$ and let $\omega \subset\subset Q$. Consider \mathcal{K} a bounded set of L^p , with $1 \leq p < \infty$. we suppose that: $\forall \epsilon > 0$, there exists $\delta > 0$, $\delta < \text{dist}(\omega, \mathbb{R}^{\ell+1} \setminus Q)$ such that $\|f(x+h) - f(x)\|_{L^p(\omega)} \leq \epsilon \forall h \in \mathbb{R}^{\ell+1}$ with $|h| < \delta$ and $\forall f \in \mathcal{K}$. Then \mathcal{K} is relatively compact in $L^p(\omega)$.

Now, we derive estimates of space and time translates of the function $\phi(u_{O, \delta t})$ which imply that the sequence is relatively compact in $L^2(Q)$.

Notice that because $(\phi(u_{O, \delta t}))_{O, \delta t}$ obey a uniform L^∞ bound, the local compactness in Q is enough to deduce the L^2 compactness.

Lemma 4.6 Let, $u_{O, \delta t}$ be the approximate solution of problem (P) defined by (11), (12). There exists a constant C_1 depending on $\Omega, T, |\phi|_{H^0}$ that

$$\int_0^T \int_{\Omega_\eta} \left| \phi(u_{O, \delta t}(t, x + \eta)) - \phi(u_{O, \delta t}(t, x)) \right|^2 dx dt \leq C_1 |\eta| \quad (36)$$

for all $\eta \in \mathbb{R}^\ell$, where $\Omega_\eta = \{x \in \Omega, [x + \eta, x] \subset \Omega\}$ and there exists C_2 depending on Ω, T, ϕ, f such that

$$\int_0^{T-\tau} \int_{\Omega} \left| \phi(u_{O, \delta t}(t + \tau, x)) - \phi(u_{O, \delta t}(t, x)) \right|^2 dx dt \leq C_2 \tau \quad (37)$$

for all $\tau \in (0, T)$.

Proof. • First, we prove (36)

Let $\eta \in \mathbb{R}^\ell$ with $\eta \neq 0$ and set $\Omega_\eta = \{x \in \Omega, [x +$

$\eta, x] \subset \Omega\}$. For all $K \in \mathcal{O}$ and $\sigma \in \varepsilon_K$, define $\chi_\sigma : \Omega_\eta \times \Omega_\eta \rightarrow \{0, 1\}$ by $\chi_\sigma(x, y) = 1$ if $[x, y] \cap \sigma \neq \emptyset$ else $\chi_\sigma(x, y) = 0$. We have

$$\begin{aligned} & \left| \phi(u_{\mathcal{O}, \delta t}(t, x + \eta)) - \phi(u_{\mathcal{O}, \delta t}(t, x)) \right| \\ & \leq \sum_{L \in \mathcal{N}(K)} \chi_{K|L}(x, x + \eta) D_{K|L} \phi; \text{ for a.e. } x \in \Omega_\eta \end{aligned} \quad (38)$$

where $D_{K|L} \phi$ is defined as

$$D_{K|L} \phi = |\phi(u_K^{n+1}) - \phi(u_L^{n+1})|.$$

We integrate (38) over Ω_η , and get:

$$\begin{aligned} & \int_{\Omega_\eta} \left| \phi(u_{\mathcal{O}, \delta t}(t, x + \eta)) - \phi(u_{\mathcal{O}, \delta t}(t, x)) \right| dx \\ & \leq \sum_{L \in \mathcal{N}(K)} \int_{\Omega_\eta} \chi_{K|L}(x, x + \eta) |D_{K|L} \phi| dx. \end{aligned} \quad (39)$$

Remark that, for all $\sigma = K|L \in \varepsilon_K$, $\int_{\Omega_\eta} \chi_{K|L}(x, x + \eta) dx$ is the measure of the set of points of Ω which are located inside the cylinder whose basis is $K|L$ and generator vector is $-\eta$. Thus

$$\int_{\Omega_\eta} \chi_{K|L}(x, x + \eta) dx \leq m(K|L) |\eta|. \quad (40)$$

The relation (39) gives

$$\begin{aligned} J &= \int_{\Omega_\eta} \left| \phi(u_{\mathcal{O}, \delta t}(t, x + \eta)) - \phi(u_{\mathcal{O}, \delta t}(t, x)) \right| dx \\ &\leq |\eta| \sum_{L \in \mathcal{N}(K)} m(K|L) |D_{K|L} \phi| \\ &\leq |\eta| \sum_{L \in \mathcal{N}(K)} m(K|L) |D_{K|L} \phi|. \end{aligned} \quad (41)$$

From now, integrate (41) over $[0, T]$

$$\int_0^T J dt \leq |\eta| \sum_{n=0}^N \delta t \sum_{L \in \mathcal{N}(K)} m(K|L) d_{K,L} \left| \frac{D_{K|L} \phi}{d_{K,L}} \right|. \quad (42)$$

Take $\Delta^\eta \phi_O^{\delta t} = \phi(u_{\mathcal{O}, \delta t}(t, x + \eta)) - \phi(u_{\mathcal{O}, \delta t}(t, x))$ and remark that:

$$\left\| \Delta \phi_O^{\delta t} \right\|_{L^2(Q_\eta)}^2 \leq \|\phi\|_{L^\infty} \left\| \Delta \phi_O^{\delta t} \right\|_{L^1(Q_\eta)}. \quad (43)$$

Then (42) and (43) give (36).

• Finally, we prove (37).

Let $\tau \in (0, T)$ and $t \in (0, t - \tau)$. Set $n_0 = [t/\delta t]$ and $n_1 = [(t + \tau)/\delta t]$, let

$$\begin{aligned} I^{t, \tau} &= \{n \in \mathbb{N}, |t < (n + 1)\delta t \leq t + \tau\} \\ J^{t, \tau} &= \{n \in \mathbb{N}, |(n + 1)\delta t - \tau \leq t < (n + 1)\delta t\}. \end{aligned}$$

Take $\Delta^\tau \phi_O^{\delta t} = \phi(u_{\mathcal{O}, \delta t}(t + \tau, x)) - \phi(u_{\mathcal{O}, \delta t}(t, x))$. Since ϕ is Locally continuous with constant $\phi_{Lip} =$

$\sup_{0 \leq a < b \leq u_{\max}} \frac{\phi(a) - \phi(b)}{a - b}$, we have:

$$\int_0^{T-\tau} \int_\Omega \left| \Delta^\tau \phi_O^{\delta t} \right|^2 dx dt \leq \phi_{Lip} \int_0^{T-\tau} S(t) dt$$

where, for almost every $t \in (0, T - \tau)$

$$\begin{aligned} & \int_\Omega \left(\Delta^\tau \phi_O^{\delta t} \right) \left(u_{\mathcal{O}, \delta t}(t + \tau, x) - u_{\mathcal{O}, \delta t}(t, x) \right) \\ &= \sum_{K \in \mathcal{O}} m(K) \left(\phi(u_K^{n_1}) - \phi(u_K^{n_0}) \right) \left(u_K^{n_1} - u_K^{n_0} \right) \\ &= \sum_{K \in \mathcal{O}} \left(\phi(u_K^{n_1}) - \phi(u_K^{n_0}) \right) \sum_{I^{t, \tau}} m(K) \left(u_K^{n+1} - u_K^n \right) \\ &= S(t). \end{aligned}$$

We get $S(t) \leq \frac{1}{2} S_0(t) + \frac{1}{2} S_1(t) + S_2(t) + S_3(t) + S_4$ with:

$$\begin{aligned} S_0(t) &= \sum_{I^{t, \tau}} \delta t \sum_{K|L} \frac{m(K|L)}{d_{K,L}} \left(\phi(u_K^{n_0}) - \phi(u_L^{n_0}) \right)^2 \\ S_1(t) &= \sum_{I^{t, \tau}} \delta t \sum_{K|L} \frac{m(K|L)}{d_{K,L}} \left(\phi(u_K^{n_1}) - \phi(u_L^{n_1}) \right)^2 \\ S_2(t) &= \sum_{I^{t, \tau}} \delta t \sum_{K|L} \frac{m(K|L)}{d_{K,L}} \left(\phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right)^2 \\ S_3(t) &= \sum_{I^{t, \tau}} \delta t \sum_{K|L} \left(\phi(u_K^{n_1}) - \phi(u_L^{n_1}) \right) F_{K,L}(u_L^{n+1}, u_K^{n+1}) \\ S_4(t) &= \sum_{I^{t, \tau}} \delta t \sum_{K|L} \left(\phi(u_L^{n_0}) - \phi(u_K^{n_0}) \right) F_{K,L}(u_L^{n+1}, u_K^{n+1}). \end{aligned}$$

We introduce the function χ^t such that $\chi^t(1) = 1$ and $\chi^t(0) = 0$. We have for all $t \in \mathbb{R}^+$ and $n \in \mathbb{N}$, $\chi^t(I^{t, \tau}) = \chi^t(J^{t, \tau})$. Therefore

$$\begin{aligned} & \int_0^{T-\tau} S_0(t) dt \\ & \leq \sum_{n=0}^{[T/\delta t]} \delta t \sum_{K|L} \frac{m(K|L)}{d_{K,L}} \left(\phi(u_K^{n_0}) - \phi(u_L^{n_0}) \right)^2 P_{n_0}^\delta \\ & \leq \sum_{n=0}^{[T/\delta t]} \delta t \sum_{K|L} m(K|L) d_{K,L} \left| \frac{\phi(u_K^{n_0}) - \phi(u_L^{n_0})}{d_{K,L}} \right|^2 P_{n_0}^\delta \\ & \leq \sum_{n=0}^{[T/\delta t]} \delta t \sum_{K|L} \ell m(\widehat{K|L}) \left| \frac{\phi(u_K^{n_0}) - \phi(u_L^{n_0})}{d_{K,L}} \right|^2 P_{n_0}^\delta. \end{aligned}$$

where

$$P_{n_0}^\delta = \int_{n_0\delta}^{(n_0+n)\delta t + \tau} \sum_{n \in \mathbb{N}} \chi^t(I^{t,\tau}) dt$$

Notice the following property:

$$\begin{aligned} & \int_{n_0\delta t}^{(n_0+1)\delta t} \sum_{n \in \mathbb{N}} \chi^t(J^{t,\tau}) dt \\ &= \sum_{n \in \mathbb{N}} \int_{(n_0-n-1)\delta t + \tau}^{(n_0-n)\delta t + \tau} \chi^t(0 \leq t < \tau) dt = \tau. \end{aligned}$$

Using (33), we find:

$$\int_0^{T-\tau} S_0(t) dt \leq C\tau. \quad (44)$$

We get in the same way

$$\int_0^{T-\tau} S_1(t) dt \leq C\tau. \quad (45)$$

We now turn to the study of the third term:

$$\begin{aligned} & \int_0^{T-\tau} S_2(t) dt \\ & \leq \sum_{n=0}^{[T/\delta t]} \delta t \sum_{K|L} \frac{m(K|L)}{d_{K,L}} \left(\phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right)^2 Q_0^\delta \\ & \leq \sum_{n=0}^{[T/\delta t]} \delta t \sum_{K|L} \ell m(\widehat{K|L}) \left| \frac{\phi(u_K^{n+1}) - \phi(u_L^{n+1})}{d_{K,L}} \right|^2 Q_{n_0}^\delta \end{aligned}$$

Because

$$\begin{aligned} & \int_0^{T-\tau} \chi^t(J^{t,\tau}) dt \\ &= \min(T - \tau, (n + 1)\delta t) - \max(0, (n + 1)\delta t - \tau) \\ &= Q_0^\delta \leq \tau, \end{aligned}$$

we get

$$\int_0^{T-\tau} S_2(t) dt \leq C\tau. \quad (46)$$

Recall that due to (10)

$$\left| \frac{F_{K,L}(a, b)}{m(K|L)} \right| \leq (\|f\|_{L^\infty} + M).$$

We have in the same way

$$\begin{aligned} & \int_0^{T-\tau} S_3(t) dt \\ & \leq \sum_{n=0}^{[T/\delta t]} \delta t \sum_{K|L} \left(\phi(u_K^{n+1}) - \phi(u_L^{n+1}) \right) F_{K,L}(u_L^{n+1}, u_K^{n+1}) Q_0^\sigma \\ & \leq \left((\|f\|_{L^\infty} + M) \sqrt{Tm(\Omega)} |\phi(u)|_{H_0} \right) \tau. \end{aligned} \quad (47)$$

In the same way we prove:

$$\begin{aligned} & \int_0^{T-\tau} S_4(t) dt \\ & \leq \left((\|f\|_{L^\infty} + M) \sqrt{Tm(\Omega)} |\phi(u)|_{H_0} \right) \tau. \end{aligned} \quad (48)$$

From (44), (45), (46), (47) and (48), we get:

$$\begin{aligned} & \int_0^{T-\tau} \int_\Omega |\phi(u_{O,\delta t}(t + \tau, x)) - \phi(u_{O,\delta t}(t, x))|^2 dx dt \\ & \leq C_2\tau. \end{aligned}$$

4.4 Existence of a discrete solution

The proof of existence of solution for the scheme (11), (12) is obtained by applying the Leray-Schauder topological degree theorem. The idea is to modify continuously the scheme to obtain a system which admits a solution and if the modification preserves in the same time the estimates (in our case this can get easily by the L^∞ norm on $u_{O,\delta t}$), then the scheme also has a solution.

Definition 4.7 Let E be a real Banach space. We denote by \mathcal{A} the set of $(Id - g, B, y)$ where $g : \bar{B} \rightarrow E$ is a compact with $B \subset E$ and $y \in E$ such that $y \notin \{g(x), x \in \partial B\}$.

Lemma 4.8 Suppose (1) is satisfied. Then for all $K \in \mathcal{O}$, there exist u_K^{n+1} satisfying (16).

Proof. For the proof, we consider for every $\alpha \in [0, 1]$ and $\forall K \in \mathcal{O}$ the following problem:

$$\begin{aligned} & v_K - \alpha \left[u_K^0 - \frac{\delta t}{m(K)} \sum_{\sigma \in \tilde{\mathcal{E}}_K} F_{K,\sigma}(v_K, v_{K,\sigma}) \right. \\ & \left. + \frac{\delta t}{m(K)} \sum_{\sigma \in \tilde{\mathcal{E}}_K} \tau_{K,\sigma} \left(\phi(v_{K,\sigma}) - \phi(v_K) \right) \right] = 0. \end{aligned}$$

with notation analogous to that of (12).

We consider the continuous function \mathcal{F} with respect to each of its variables defined by:

$$\begin{aligned} \mathcal{F}(\alpha, v) &= v_K - \alpha \left[u_K^0 - \frac{\delta t}{m(K)} \sum_{\sigma \in \tilde{\mathcal{E}}_K} F_{K,\sigma}(v_K, v_{K,\sigma}) \right. \\ & \left. + \frac{\delta t}{m(K)} \sum_{\sigma \in \tilde{\mathcal{E}}_K} \tau_{K,\sigma} \left(\phi(v_{K,\sigma}) - \phi(v_K) \right) \right]. \end{aligned}$$

The function $\mathcal{F}(\alpha, \cdot)$ is a continuous homotopy between $\mathcal{F}(0, \cdot)$ and $\mathcal{F}(1, \cdot)$. First, remark that $u_K^{n+1} = 0$ is solution of $\mathcal{F}(0, u_K^{n+1}) = 0$ for all $(n, K) \in [0, N] \times \mathcal{O}$. If B is a ball with a sufficiently large radius in the space of solution of the system, the equation

$\mathcal{F}(\cdot, \cdot) = 0$ has no solution on the boundary ∂B . Indeed replacing u_0, f, ϕ by $\alpha u_0, \alpha f, \alpha \phi$ we can apply the argument of Proposition 4.1 to solutions of equation $\mathcal{F}(\alpha, v) = 0$. Then it is enough to supply the finite dimensional set \mathbb{R}^θ of discrete functions by the norm $\|\cdot\|_{L^\infty}$ and take B of radius larger than u_{\max} . Therefore

$$\text{degree}(\mathcal{F}(0, \cdot), B) = \text{degree}(\mathcal{F}(1, \cdot), B) \neq 0.$$

Thus there exists at least a solution to equation $\mathcal{F}(1, \cdot) = 0$. This solution is a solution to our scheme.

5 Continuous entropy inequality

We prove in this section that the approximate solutions fulfil a continuous entropy inequality in the sense of Theorem 5.2 below. Before, we recall a result that will serve us in the proof of this Theorem.

Lemma 5.1 *e.g., [12] Let K be a non empty open convex polygonal set in \mathbb{R}^ℓ . For $\sigma \in \bar{\varepsilon}_K$, we denote by x_σ the center of gravity of σ ; we also denote by $n_{K,\sigma}$ the unit normal vector to σ outward to K . Then, for all vector $\vec{V} \in \mathbb{R}^\ell$ and for all point $x_K \in K$, we have:*

$$m(K)\vec{V} = \sum_{\sigma \in \bar{\varepsilon}_K} m(\sigma)\vec{V} \cdot n_{K,\sigma}(x_\sigma - x_K). \quad (49)$$

Proof. We denote by a superscript i , the i -th coordinate of vectors and points in \mathbb{R}^ℓ . By Stokes formula, we have:

$$\begin{aligned} m(K)V^i &= \int_K \text{div}((x^i - x_K^i)\vec{V}) dx \\ &= \int_{\partial K} (x^i - x_K^i)\vec{V} \cdot n_K d\gamma(x) \\ &= \sum_{\sigma \in \bar{\varepsilon}_K} \int_\sigma (x^i - x_K^i)\vec{V} \cdot n_{K,\sigma} d\gamma(x). \end{aligned} \quad (50)$$

Hence, by the definition of the center of gravity, we have:

$$\begin{aligned} \int_\sigma (x^i - x_K^i) d\gamma(x) &= \int_\sigma x^i d\gamma(x) - m(\sigma)x_K^i \\ &= m(\sigma)x_\sigma^i - m(\sigma)x_K^i. \end{aligned} \quad (51)$$

Replace (51) in (50); we find (49).

From now, as the approximate solutions satisfy the discrete entropy inequalities (22), we prove that they satisfy a continuous form of these inequalities.

Theorem 5.2 *Assume that (3), (6)-(9) hold. Let $u_{O,\delta t}$ be the approximate solution of the problem (P) defined by (11),(12). Then the following continuous*

approximate entropy inequalities hold: for all $k \in [0, u_{\max}]$, for all $\xi \in C^\infty([0, T] \times \mathbb{R}^\ell)$, $\xi \geq 0$,

$$\begin{aligned} &\int_0^T \int_\Omega \eta_k(u_{O,\delta t}) \xi_t dx dt + \int_\Omega \eta_k(u_0) \xi(0, x) dx \\ &+ \int_0^T \int_\Omega \left(\Phi_k(u_{O,\delta t}) - \nabla_O \eta_{\phi(k)}(\phi(u_{O,\delta t})) \right) \cdot \nabla \xi dx dt \\ &+ \int_0^T \int_{\partial\Omega} |f(k) \cdot \eta(x)| \xi(t, x) d\mathcal{H}^{\ell-1}(x) dt \\ &\geq -v_{O,n}(\xi); \end{aligned} \quad (52)$$

where: $\forall \xi \in C^\infty([0, T] \times \mathbb{R}^\ell)$, $v_{O,n}(\xi) \rightarrow 0$ when $h \rightarrow 0$. Here

$$\begin{aligned} &\sum_{n=0}^N \sum_{K|L} 1_{K|L \times [t_n, t_{n+1}]} \nabla_{K|L} \eta_{\phi(k)}(\phi(u_{O,\delta t})) \\ &= \nabla_O \eta_{\phi(k)}(\phi(u_{O,\delta t})). \end{aligned}$$

Remark 5.3 *In the same case, if we replace in (52) η_k by η_k^+ (resp η_k^-) and $|f(k) \cdot \eta(x)|$ by $(f(k) \cdot \eta(x))^+$ (resp $(f(k) \cdot \eta(x))^-$) we obtain sub entropy inequalities (resp super entropy inequalities). Obviously, the approximate solution $u_{O,\delta t}$ is an approximate entropy solution if and only if $u_{O,\delta t}$ is approximate entropy sub-solution and entropy super-solution simultaneously.*

Proof of Theorem 5.2 Let $\xi \in C^\infty([0, T] \times \mathbb{R}^\ell)^+$ and $k \in [0, u_{\max}]$, we fix $T \geq 0$ and set $N = \frac{T}{\delta t} + 1$. It is enough to suppose that $\xi(t, x) = \theta(t)\zeta(x)$, this mean that $\xi_K^{n+1} = \theta^{n+1}\zeta_K$. By density in $C^\infty([0, T] \times \mathbb{R}^\ell)$ of linear combinations of such functions, the general case will follow. Depending on the circumstances, $\zeta_K = \int_K \zeta$ or $\zeta_K = \zeta(x_K)$ with x_K the center of control volume K . Take $\tilde{\zeta}_K = \int_K \zeta(x) dx - \zeta_K$. Multiplying inequality (22) by $\delta t \xi_K^{n+1}$ and summing over $K \in \mathcal{O}$ and $n \in \{0, \dots, N\}$, yields the inequality

$$\begin{aligned} &\sum_{n=0}^N \sum_{K \in \mathcal{O}} m(K) \left(\eta_k(u_K^{n+1}) - \eta_k(u_K^n) \right) \xi_K^{n+1} \\ &+ \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} \sum_{\sigma \in \varepsilon_K} \Phi_{K,\sigma,k}(u_K^{n+1}, u_{K,\sigma}^{n+1}) \xi_K^{n+1} \\ &- \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} \sum_{\sigma \in \varepsilon_K^{ext}} \text{sign}(u_K^{n+1} - k) F_{K,\sigma,k}(k, k) \xi_K^{n+1} \\ &- \sum_{n=0}^N \delta t \sum_{K|L} \sum_{K|L} \tau_{K|L} \Delta \eta_{\phi(k)}^{K,L} \phi \xi_K^{n+1} \leq 0. \end{aligned} \quad (53)$$

with $\Delta \eta_{\phi(k)}^{K,L} \phi = \eta_{\phi(k)}(\phi(u_L^{n+1})) - \eta_{\phi(k)}(\phi(u_K^{n+1}))$. Hence (53) can be seen as $I_{E_v}^d + I_{C_v}^d + I_{Dif}^d \leq 0$.

To prove inequality (52), we have to prove that $I_{E_v}^c + I_{C_v}^c + I_{Dif}^c \leq \nu_{O,n}(\xi)$ where $I_{E_v}^c$, $I_{C_v}^c$ and I_{Dif}^c are defined by:

$$I_{E_v}^c = - \int_0^T \int_{\Omega} \eta_k(u_{O,\delta t}) \zeta(x) \theta_t(t) dx dt - \int_{\Omega} \eta_k(u_0) \theta(0) \zeta(x) dx,$$

$$I_{C_v}^c = - \int_0^T \theta \int_{\Omega} \Phi_k(u_{O,\delta t}) \cdot \nabla \zeta dx dt - \int_0^T \theta \int_{\partial\Omega} |f(k) \cdot \eta(x)| \zeta(x) d\mathcal{H}^{\ell-1}(x) dt,$$

$$I_{Dif}^c = \int_0^T \theta \int_{\Omega} \nabla_O \eta_{\phi(k)}(\phi(u_{O,\delta t})) \cdot \nabla \zeta dx dt.$$

Then, we have to compare $I_{E_v}^d$ with $I_{E_v}^c$; $I_{C_v}^d$ with $I_{C_v}^c$; and I_{Dif}^d with I_{Dif}^c .

Firstly, we have to estimate $|I_{E_v}^d - I_{E_v}^c|$. Using the definition of $u_{O,\delta t}$, the quantity $I_{E_v}^d$ reads:

$$\begin{aligned} I_{E_v}^d &= - \sum_{n=0}^{N-1} \sum_{K \in O} m(K) \eta_k(u_K^{n+1}) \left(\xi_K^{n+1} - \xi_K^n \right) \\ &\quad - \sum_{K \in O} m(K) \left(\eta_k(u_K^0) \xi_K^1 - \eta_k(u_K^{N+1}) \xi_K^{N+1} \right) \\ &= - \sum_{n=0}^{N-1} \delta t \sum_{K \in O} m(K) \eta_k(u_K^{n+1}) \frac{\xi_K^{n+1} - \xi_K^n}{\delta t} \\ &\quad - \sum_{K \in O} m(K) \eta_k(u_K^0) \xi_K^1 \\ &= - \sum_{n=0}^{N-1} \delta t \sum_{K \in O} m(K) \eta_k(u_K^{n+1}) \frac{\theta^{n+1} - \theta^n}{\delta t} \zeta_K \\ &\quad - \sum_{K \in O} m(K) \eta_k(u_K^0) \theta^1 \zeta_K \\ &= - \sum_{n=0}^{N-1} \delta t \sum_{K \in O} m(K) \eta_k(u_K^{n+1}) (\theta^n)_t \int_K \zeta(x) dx \\ &\quad - \sum_{K \in O} m(K) \eta_k(u_K^0) \theta^1 \int_K \zeta(x) dx, \end{aligned} \tag{54}$$

with $(\theta^n)_t = \int_{t_n}^{t_{n+1}} \theta_t dt$. We deduce that

$$|I_{E_v}^d - I_{E_v}^c| \leq \nu_{O,k}^1(\xi) + \nu_{O,k}^2(\xi), \tag{55}$$

where:

$$\sum_{n=1}^{N-1} \delta t \sum_{K \in O} m(K) \eta_k(u_{O,\delta t}) \mathbf{1}_{K \times [t_n, t_{n+1}]}(\theta^n)_t |\zeta_K| = \nu_{O,k}^1(\xi) \tag{56}$$

$$\nu_{O,k}^2(\xi) = \sum_{K \in O} m(K) \eta_k(u_K^0) \theta^1 |\zeta_K|. \tag{57}$$

As $\xi \in C^\infty$, then we have:

$$|\zeta_K| = \left| \int_K \zeta(x) dx - \zeta_K \right| \leq \|\zeta\|_{C^1} h.$$

Then, the quantities $\nu_{O,k}^1(\xi)$, $\nu_{O,k}^2(\xi)$, tend to zero when $h \rightarrow 0$.

Secondly, we study the difference between $I_{C_v}^d$ and $I_{C_v}^c$. We take care separately of what happens inside and what happens on the boundary of Ω . Therefore we write $I_{C_v}^c$ has the sum of $I_{C_v}^{c,int}$ and $I_{C_v}^{c,ext}$.

$$I_{C_v}^{c,int} = - \int_0^T \theta \int_{\Omega} \Phi_k(u_{O,\delta t}) \cdot \nabla \zeta dx dt$$

$$I_{C_v}^{c,ext} = - \int_0^T \theta \int_{\partial\Omega} |f(k) \cdot \eta(x)| \zeta(x) d\mathcal{H}^{\ell-1}(x) dt.$$

Further, introduce auxiliary values $(\zeta_{K|L})_{L \in \mathcal{N}(K)}$ by $\zeta_{K|L} = \zeta(x_{K|L})$, where $x_{K|L}$ is the barycenter of $K|L$. Denote by $\Delta_K \xi^{n+1} = \xi_K^{n+1} - \xi_{K|L}^{n+1}$. The term $I_{C_v}^d$, which can be rewritten as the sum between $I_{C_v}^{d,int}$ and $I_{C_v}^{d,ext}$:

$$\sum_{n=0}^N \delta t \sum_{\sigma \in K|L} \Phi_{K,K|L,k}(u_K^{n+1}, u_L^{n+1}) [\Delta_K \xi^{n+1} - \Delta^L \xi^{n+1}]$$

$$= \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{L \in \mathcal{N}(K)} \Phi_{K,K|L,k}(u_K^{n+1}, u_L^{n+1}) \Delta^K \xi^{n+1}$$

$$= \sum_{n=0}^N \theta^{n+1} \delta t \sum_{K \in O} \sum_{L \in \mathcal{N}(K)} \Phi_{K,K|L,k}(u_K^{n+1}, u_L^{n+1}) \Delta^K \zeta$$

$$= I_{C_v}^{d,int}$$

$$- \sum_{n=0}^N \delta t \sum_{K \in O} \sum_{\sigma \in \mathcal{E}_K^{ext}} \text{sign}(u_K^{n+1} - k) F_{K,\sigma}(k, k) \xi_K^{n+1}$$

$$= - \sum_{n=0}^N \theta^{n+1} \delta t \sum_{K \in O} \sum_{\sigma \in \mathcal{E}_K^{ext}} \text{sign}(u_K^{n+1} - k) F_{K,\sigma}(k, k) \zeta_K$$

$$= I_{C_v}^{d,ext}.$$

Now, we compare $I_{Cv}^{c,int}$ and $I_{Cv}^{d,int}$. As the numerical fluxes, the numerical entropy fluxes are consistent:

$$\begin{aligned} & \int_{\sigma} \Phi_k(u_K^{n+1}).n_{K,\sigma}d\gamma(x)dt \\ &= m(K|L)\Phi_k(u_K^{n+1}).n_{K|L} \\ &= \Phi_{K,\sigma,k}(u_K^{n+1}, u_K^{n+1}). \end{aligned} \quad (58)$$

Simultaneously, for each $K \in \mathcal{O}$, we approach ζ by the affine function $\tilde{\zeta}_K$ in a neighbourhood of K , with $\tilde{\zeta}_K = \tilde{\zeta}(x_K)$, we set $\tilde{\zeta}_{K|L} = \tilde{\zeta}(x_{K|L})$. Then

$$\begin{aligned} \zeta(x)1_K &= \tilde{\zeta}_K + \underline{o}(|x - x_K|^2); \\ \zeta_{K|L} - \tilde{\zeta}_{K|L} &= \underline{o}(h^2); \\ \nabla \tilde{\zeta}_K &= \text{cst on } K \\ \|\nabla \zeta - \nabla \tilde{\zeta}\|_{L^\infty(K)} &= \underline{o}(h) \text{ and} \\ \nabla \tilde{\zeta}_K.(x_K - x_{K|L}) &= \tilde{\zeta}_K - \tilde{\zeta}_{K|L}. \end{aligned} \quad (59)$$

We denote the resulting expression by $\tilde{I}_{Cv}^{d,int}$, we have

$$\begin{aligned} & - \sum_{n=0}^N \delta t \sum_K \sum_L \Phi_{K,K|L,k}(u_K^{n+1}, u_K^{n+1})(-\Delta_K \tilde{\xi}^{n+1}) \\ &= \tilde{I}_{Cv}^{d,int} \end{aligned}$$

From now, using (58) and Lemma 5.1, which states that

$$\sum_{K|L} m(K|L)\nabla \tilde{\zeta}_K.n_{K|L}(x_{K|L} - x_K) = m(K)\nabla \tilde{\zeta}_K,$$

we find:

$$\tilde{I}_{Cv}^{d,int} = - \sum_{n=0}^N \delta t \theta^{n+1} \sum_{K \in \mathcal{O}} \Phi_k(u_K^{n+1})m(K)\nabla \tilde{\zeta}_K.$$

It is easy to see that

$$\tilde{I}_{Cv}^{d,int} = - \int_0^T \theta \int_{\Omega} \Phi_k(u_{\mathcal{O},\delta t}).\nabla \tilde{\zeta}_K dxdt =: \tilde{I}_{Cv}^{d,int}$$

$$\begin{aligned} |I_{Cv}^{d,int} - I_{Cv}^{c,int}| &\leq |I_{Cv}^{d,int} - \tilde{I}_{Cv}^{d,int}| + |\tilde{I}_{Cv}^{d,int} - I_{Cv}^{c,int}| \\ &= v_{\mathcal{O},k}^3(\xi) + v_{\mathcal{O},k}^4(\xi) \end{aligned}$$

with:

$$v_{\mathcal{O},k}^3(\xi) = \sum_{n=0}^N \delta t \theta^{n+1} \sum_{K|L} \left| \Delta_{K,L}^k \overline{\Phi_{K,L,k}} \Delta_K \zeta \right|; \quad (60)$$

$$v_{\mathcal{O},k}^4(\xi) = \int_0^T \theta \int_{\Omega} \Phi_k(u_{\mathcal{O},\delta t}).|\nabla \zeta - \nabla \tilde{\zeta}_K| dxdt. \quad (61)$$

where

$$\begin{aligned} & \Phi_{K,L,k}(u_K^{n+1}, u_K^{n+1}) - \Phi_{K,L,k}(u_K^{n+1}, u_L^{n+1}) \\ &= \Delta_{K,L}^k \overline{\Phi_{K,L,k}}. \end{aligned}$$

Let us show that $v_{\mathcal{O},k}^3(\xi)$ and $v_{\mathcal{O},k}^4(\xi)$ tend to zero as $h \rightarrow 0$. Thanks to (59), $v_{\mathcal{O},k}^4(\xi)$ as $h \rightarrow 0$. Now, we write:

$$\begin{aligned} & \frac{1}{\delta t m(K|L)} \int_{n\delta t}^{(n+1)\delta t} \int_{K|L} (\xi(t,x) - \xi(s,y))d\gamma(y)ds \\ &= \xi(t,x) - \hat{\xi}_{K|L}^{n+1}. \end{aligned} \quad (62)$$

For all $(x,y) \in K|L \times K|L$,

$$|\zeta(x) - \zeta(y)| \leq h \|\nabla \zeta\|_{L^\infty}. \quad (63)$$

We exploit the BV-weak estimates on space derivatives to prove that $v_{\mathcal{O},k}^3(\xi)$ tend to zero when h goes to zero. Indeed, we have

$$|\Delta_{K,L}^k \overline{\Phi_{K,L,k}}| \leq \max_{i_\alpha} (F_{K,\sigma}(d,c) - (F_{K,\sigma}(d,d)))$$

and thanks to (63), we get an estimate on the difference between the average value of ζ and a control volume and on one of its edges: there exists C_ζ depending only upon ζ , such that

$$\forall K|L, |\zeta_K - \zeta_{K|L}^{n+1}| \leq C_\zeta h.$$

Therefore, the following estimate on $v_{\mathcal{O},k}^3(\xi)$ holds:

$$\begin{aligned} C_\zeta(h) & \sum_{n=0}^N \delta t \sum_{K|L} \left[\max_{i_\alpha} (F_{K,\sigma}(d,c) - F_{K,\sigma}(d,d)) \right] \\ & + C_\zeta(h) \sum_{n=0}^N \delta t \sum_{K|L} \left[\max_{i_\alpha} (F_{K,\sigma}(d,c) - F_{K,\sigma}(c,c)) \right] \\ &= v_{\mathcal{O},k}^3(\zeta) \leq C_\zeta \frac{h}{\sqrt{h}} \end{aligned}$$

where the constant C_ζ is given by (25). Now, it remains to notice that

$$\begin{aligned} & \sum_{n=0}^N \theta^{n+1} \delta t \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K^{ext}} \text{sign}(u_K^{n+1} - k) F_{K,\sigma}(k,k) \zeta_K \\ &= -I_{Cv}^{d,ext} \\ &\leq \sum_{n=0}^N \theta^{n+1} \delta t \sum_{K \in \mathcal{O}} \sum_{\sigma \in \mathcal{E}_K^{ext}} |F_{K,\sigma}(k,k)| \zeta_K = -I_{Cv}^{c,ext}. \end{aligned}$$

Then, we have:

$$I_{Cv}^{c,ext} - I_{Conv}^{d,ext} \leq 0.$$

The last step is to compare $I_{Dif}^{c,ext}$ to $I_{Dif}^{d,ext}$. We rewrite the term I_{Dif}^d as

$$\begin{aligned} & - \sum_{n=0}^N \delta t \sum_{\sigma \in K|L} \tau_{K|L} \Delta \eta_{\phi(k)}^{K,L} \left(\xi_K^{n+1} - \xi_L^{n+1} \right) \\ & = - \sum_{n=0}^N \delta t \theta^{n+1} \sum_{\sigma \in K|L} \frac{m(K|L)}{d_{K,L}} \Delta \eta_{\phi(k)}^{K,L} \left(\zeta_K - \zeta_L \right) \\ & = - \sum_{n=0}^N \delta t \theta^{n+1} \sum_{\sigma \in K|L} m(\widehat{K|L}) \nabla_{\widehat{K|L}} \eta_{\phi(k)} (\phi(u_O^{n+1})) \tilde{\nabla}_{\widehat{K|L}} \zeta \\ & = I_{Dif}^d \end{aligned}$$

where: $\tilde{\nabla}_{\widehat{K|L}} \zeta = \int_{x_K}^{x_L} \nabla \zeta$. Notice that

$$\|\nabla \zeta - \tilde{\nabla}_{\widehat{K|L}} \zeta\|_{L^\infty(\widehat{K|L})} = \underline{o}(h).$$

Therefore we have

$$|I_{Dif}^d - I_{Dif}^c| \leq \nu_{O,k}^5(\xi), \quad (64)$$

with:

$$\begin{aligned} & \int_0^T \int_{\Omega} |\nabla_O \eta_{\phi(k)}(\phi(u_{O,\delta t}))| \cdot |\nabla \zeta - \tilde{\nabla}_{\widehat{K|L}} \zeta| dx dt \\ & = \nu_{O,k}^5(\xi). \end{aligned} \quad (65)$$

To conclude, we prove that $\nu_{O,k}^5(\xi) \rightarrow 0$ as $h \rightarrow 0$. Using Cauchy-Schwartz inequality, we find

$$\nu_{O,k}^5(\xi) \leq \|\theta\|_{L^\infty} \|\nabla_O \eta_{\phi(k)}(\phi(u_{O,\delta t}))\|_{L^2} \underline{o}(h).$$

Then, using the fact that η is 1-Lipschitz, and the estimate (33) we prove that $\nu_{O,k}^5(\xi) \rightarrow 0$ as $h \rightarrow 0$.

6 Convergence of the scheme

The main result of this paper is the following theorem.

Theorem 6.1 (Convergence of the approximate solution towards the entropy solution). Assume that one of the following hypotheses is satisfied

$$\ell = 1 \text{ and } \Omega = (a, b) \text{ an interval of } \mathbb{R}; \quad (66)$$

$$\ell \geq 1 \text{ } u_c = 0, \text{ and } f \circ \phi^{-1} \in C^{0,\alpha}, \alpha > 0; \quad (67)$$

$$\ell \geq 1 \text{ } u_c = u_{\max}. \quad (68)$$

Let, $(u_{O,\delta t})_{O,\delta t}$ be a family of approximate solutions of problem (P) defined by (11), (12). Then, under hypotheses (4)-(9), we have $\max(\delta t, h) \rightarrow 0$:

$$\forall p \in [1, +\infty) \text{ } u_{O,\delta t} \rightarrow u \text{ in } L^p(Q) \quad (69)$$

$$\nabla_O \phi(u_{O,n}) \rightarrow \nabla \phi(u) \text{ in } L^2(Q)$$

where u is the unique entropy solution of (P), i.e u satisfies (2).

Remark 6.2 It is possible to replace in the Theorem 6.1 all the three hypotheses (66), (67), (68) by the following one, which is much more general:

$$(H) \left\{ \begin{array}{l} \ell \geq 1 \text{ and } u_0 \text{ is such that there exist an} \\ \text{entropy solution of (P) such that} \\ (f(u) - \nabla \phi(u)) \cdot \eta(x) \\ \text{possess a strong trace in } L^1 \text{ sense.} \end{array} \right.$$

Such kind of functions u which verify (H), will be called trace regular entropy solutions in reference [13], [14]. The idea to prove uniqueness of entropy solution is to compare any entropy solution of (P) with trace regular entropy solution and break the symmetry in the application of doubling of variables method by taking test function that is zero on the boundary $\bar{Q} \times ((0, T) \times \partial \Omega)$ of $Q \times Q$ but non zero on the boundary $((0, T) \times \partial \Omega) \times \bar{Q}$ e.g. the method of [1], [6]. If (H) is satisfied for all u_0 that belong to a certain subset X such that $\bar{X}^{\|\cdot\|_{L^1}} = L^1(\Omega; [0, u_{\max}])$, then uniqueness is true for all u_0 .

Presently to our knowledge the only results which establish that (H) holds for a dense subset X is proved for the case (68), [4], [15].

In this pure hyperbolic case existence of the strong trace of the flux is established in [4], [15]. Then uniqueness of entropy solution follows by standard doubling of variables method and it is enough to take a symmetric test function.

In the case where hypotheses (67) or (66) are satisfied, it is more easy to prove existence of trace regular entropy solution for the stationary problem with L^∞ source term. In this case, we even have sense that the total flux is continuous up to the boundary, i.e $(f(u) - \nabla \phi(u)) \cdot \eta \in C(\bar{\Omega})$ for example [16], [17]. Then we can adopt the same strategy as in the case where (H) holds, but in the doubling of variables method we compare entropy solution of (P) with trace regular entropy solution of (S). Then using nonlinear semigroup approach, we proved that entropy solution of (P) is the unique mild solution, the reader can refer to [1], [6], [18]. The same strategy is adopted here to prove that entropy-process solution (Definition 6.3) is the unique entropy solution (see Appendix 1 and 2).

Proof of Theorem 6.1 The proof of Theorem 6.1 is in two steps. First in Proposition 6.8, we prove that the approximate solutions converge towards an entropy-process solution. Then in Appendix 2 (Theorem 8.4, and Proposition 8.5, 8.6, 8.7) we prove that entropy-process solution is in fact the unique entropy solution using the intermediate notion of integral-process solution developed for this purpose in the Appendix 1.

6.1 Entropy process solution

Definition 6.3 Let $\mu \in L^\infty(Q \times (0, 1))$. The function $\mu = \mu(t, x, \alpha)$ taking values in $[0, u_{\max}]$ is called an entropy-process solution to problem (P) if $\forall k \in [0, u_{\max}], \forall \xi \in C^\infty([0, T] \times \mathbb{R}^\ell)$, with $\xi \geq 0$, the following inequality holds :

$$\begin{aligned} & \int_0^T \int_\Omega \int_0^1 |\mu(\alpha)u - k|\xi_t dx dt d\alpha + \int_\Omega |u_0 - k|\xi(0, x) dx \\ & + \int_0^T \int_\Omega \int_0^1 \text{sign}(\mu(\alpha) - k) [f(\mu) - f(k)] \cdot \nabla \xi dx dt d\alpha \\ & - \int_0^T \int_\Omega \nabla |\phi(u) - \phi(k)| \cdot \nabla \xi dx dt \\ & + \int_0^T \int_{\partial\Omega} |f(k) \cdot \eta(x)| \xi(t, x) d\mathcal{H}^{\ell-1}(x) dt \geq 0, \quad (70) \end{aligned}$$

where $u(t, x) = \int_0^1 \mu(t, x, \alpha) d\alpha$.

Remark 6.4 If $\mu \in L^\infty(Q \times (0, 1))$ is entropy process solution then, it satisfies for all $\xi \in L^2(0, T; H^1(\Omega))$ such that $\xi_t \in L^1(Q)$ and $\xi(T, \cdot) = 0$

$$\begin{aligned} & \int_0^1 \int_0^T \int_\Omega \left\{ \mu \xi_t + \left(f(\mu) - \nabla \phi(u) \right) \cdot \nabla \xi \right\} dx dt d\alpha \\ & + \int_\Omega u_0 \xi(0, x) dx = 0. \quad (71) \end{aligned}$$

We recall the nonlinear weak star convergence for $(u_{O, \delta t})_{O, \delta t}$ which is equivalent to the notion of convergence towards a Young measure as developed in [10].

Theorem 6.5 ([19]) (Nonlinear weak star Convergence) Let $(u_n)_{n \in \mathbb{N}}$ be a bounded sequence in $L^\infty(Q)$. Then, there exists $\mu \in L^\infty(Q \times (0, 1))$, such that up to a subsequence, u_n tends to μ in the nonlinear weak star sense as $n \rightarrow \infty$, i.e: $\forall h \in C(\mathbb{R}, \mathbb{R})$

$$h(u_n) \rightharpoonup \int_0^1 h(\mu(\cdot, \alpha)) d\alpha \text{ weakly } - * \text{ in } L^\infty(Q) \quad (72)$$

Moreover, if μ is independent on α (i.e $\mu(t, x, \alpha) = u(t, x)$ for a.e. (t, x) , and for all α), then u_n converge strongly in $L^1(Q)$ towards some $u(t, x)$. In particular, observe that the following holds:

Lemma 6.6 Suppose that the sequence $u_n(\cdot) \rightarrow \mu(\cdot, \alpha)$ in the nonlinear weak star sense, assume that g is a continuous non decreasing function such that $g(u_n(\cdot)) \rightarrow \theta$ strongly in $L^1(Q)$. Then, $\theta = g(\mu(\cdot, \alpha)) = g(u)$ where $u(t, x) = \int_0^1 \mu(t, x, \alpha) d\alpha$.

Proof. Let $v_n = g(u_n)$, since g is continuous, then the sequence v_n is bounded in $L^\infty(Q)$, so that $v_n(t, x) \xrightarrow{nl-*} v(t, x, \alpha)$ (where $\xrightarrow{nl-*}$ mean the convergence for weak star topology in $L^\infty(Q)$) and $v_n \rightarrow \theta$ in $L^1(Q)$ and $v(t, x, \alpha) := g(\mu(t, x, \alpha))$ is an associated Young measure, since for all $h \in C(\mathbb{R}, \mathbb{R})$

$$\begin{aligned} h(v_n(t, x)) &= (h \circ g)(u_n) \rightharpoonup \int_0^1 h \circ g(\mu(\cdot, \alpha)) d\alpha \\ &= \int_0^1 h(v(t, x, \alpha)) d\alpha. \end{aligned}$$

Since v_n tend to θ strongly, we deduce that $v(t, x, \alpha) = \theta(t, x)$ and v does not depend on α . Moreover, if g is continuous and nondecreasing the level sets $g^{-1}(\{c\})$ are closed intervals of \mathbb{R} . Then for all $(t, x) \in Q$,

$$\begin{aligned} \mu(\cdot, \alpha) \in g^{-1}(\{\theta(\cdot)\}) &\Rightarrow u(\cdot) = \int \mu(\cdot, \alpha) d\alpha \\ \int \mu(\cdot, \alpha) d\alpha \in g^{-1}(\{\theta(\cdot)\}) &\Rightarrow g(u(\cdot)) = \theta(\cdot). \end{aligned}$$

From now we give a "discrete $L^2(0, T; H^1(\Omega))$ " compactness result e.g., [19], [20], [21], [22], [23].

6.2 Convergence towards an entropy-process solution

Lemma 6.7 Consider a family of corresponding discrete functions $w_{O, \delta t}$ satisfying the uniform bounds.

$$\begin{aligned} \sum_{n=0}^N \delta t \sum_{K \in \mathcal{O}} m(K) (w_K^{N+1})^2 &\leq C; \\ \sum_{n=0}^N \delta t \sum_{K|L} \tau_{K|L} (\nabla_{\overline{K|L}} w_O)^2 &\leq C, \end{aligned}$$

where the discrete gradient $\nabla_{\overline{K|L}}$ are defined by (5).

Then there exists $w \in L^2(0, T; H^1(\Omega))$ such that, up to extraction of a subsequence, $w_{O, \delta t} \rightarrow w$ in $L^2(Q)$ weakly and $\nabla_O w \rightarrow \nabla w$ in $(L^2(Q))^\ell$ weakly.

We wish to prove the convergence of the approximate solution $(u_{O, \delta t})$ to an entropy solution u of (P), i.e. we want to prove that there exists a limit u and that it satisfies (2). For this purpose, we prove first that $(u_{O, \delta t})$ tends in the nonlinear weak star sense to an entropy-process solution.

Proposition 6.8 (Convergence towards an entropy-process solution) Under hypotheses (4)-(9), let $u_{O, \delta t}$ be the approximate solution of problem (P) defined by (11), (12). There exists an entropy-process solution μ of (P) in the sense of Definition 6.3 and a subsequence of $(u_{O, \delta t})_{O, \delta t}$, such that:

1. The sequence $(u_{O,\delta t})_{O,\delta t}$ converges to μ in the nonlinear weak star sense.
2. Moreover $(\phi(u_{O,\delta t}))_{O,\delta t}$ converges strongly in $L^2(Q)$ to $\phi(u)$ as $h, \delta t$ tend to zero and
3. $(\nabla_O \phi(u_{O,\delta t}))_{O,\delta t} \rightharpoonup \nabla \phi(u)$ in $(L^2(Q))^\ell$ weakly,

where $u(t, x) = \int_0^1 \mu(t, x, \alpha) d\alpha$.

From this result, we deduce Theorem 6.1, using additional regularity properties coming from (66), (67) or (68) also Remark 6.2 for variants of the concluding argument.

Proof of Proposition 6.8 Passage to the limit in the continuous entropy inequality:

Recall that we have proved that $v_{O,n}(\xi) \rightarrow 0$ when $(h, \delta t) \rightarrow 0$ for $\xi \in C^\infty([0, T] \times \mathbb{R}^\ell)$. We follow step by step the passage to the limit for each term of the left hand side of (5.2).

Because $u_{O,\delta t}$ is bounded in $L^\infty(Q)$, by Theorem 6.5, there exist $\mu \in L^\infty(Q \times (0, 1))$ such that up to a subsequence, $(u_{O,\delta t})$ tends to μ in the nonlinear weak star sense as $\max(\delta t, h) \rightarrow 0$. We set $u(t, x) = \int_0^1 \mu(t, x, \alpha) d\alpha$. Using the continuity of $\Phi_k(\cdot)$ and $\eta_k(\cdot) = |\cdot - k|$, we prove that:

$$\int_0^T \int_\Omega \eta_k(u_{O,\delta t}) \xi_t \rightarrow \int_0^T \int_\Omega \int_0^1 \mu(t, x, \alpha) \xi_t dx dt d\alpha,$$

$$\int_0^T \int_\Omega \Phi_k(u_{O,\delta t}) \cdot \nabla \xi dx dt \rightarrow \int_0^T \int_\Omega \int_0^1 \Phi_k(\mu) dx dt d\alpha.$$

Due to (33) and by the Fréchet-Kolmogorov's theorem (due to the time and space translation on $\phi(u_{O,\delta t})$) we can apply lemma 6.7 for $w_{O,\delta t} = \phi(u_{O,\delta t})$. Notice that in view of Lemma 6.6, it appears that $\phi(\mu) = \phi(u)$ where $u(t, x) = \int_0^1 \mu(t, x, \alpha) d\alpha$. The Lipschitz continuity of η_ϕ permits to have

$$\int_0^T \int_\Omega \nabla_O \eta_\phi(\phi(u_{O,n})) \cdot \nabla \xi \rightarrow \int_0^T \int_\Omega \nabla \eta_\phi(\phi(u)) \cdot \nabla \xi.$$

We conclude that $u_{O,\delta t}$ converge to an entropy-process solution μ .

Acknowledgment

I would like to thank Boris Andreianov and Petra Wittbold for their helpful readings and remarks that helped me improve this article. I would like to thank the anonymous referees for their careful reading of the article and their very useful remarks.

References:

- [1] Andreianov, B. and Karimou Gazibo, M., Entropy Formulation of Degenerate Parabolic Equation with Zero-Flux Boundary Condition. *Zeitschrift für Angewandte Mathematik und Physik*, **64**, (2013), 1471-1491. <https://doi.org/10.1007/s00033-012-0297-6>
- [2] Carrillo, J. Entropy Solutions for Nonlinear Degenerate Problems, *Archive for Rational Mechanics and Analysis*, **147**, No 4, (1999), 269-361. <https://doi.org/10.1007/s002050050152>
- [3] Kruzkov, S.N., First order quasi-linear equations in several independent variables, *Matematicheskii Sbornik*, **10**, (2), (1970), 217-243.
- [4] R. Bürger, Frid, H. and Karlsen, K.H. On the Well-Posedness of Entropy Solutions to Conservation Laws with a Zero-Flux Boundary Condition, *Journal of Mathematical Analysis and Applications*, **326**, (2007), 108-120. <https://doi.org/10.1016/j.jmaa.2006.02.072>.
- [5] K.H. Karlsen and J.D. Towers, Convergence of monotone schemes for conservation laws with zero-flux boundary conditions, *Advances in Applied Mathematics and Mechanics*, **9**, No.3, (2017), 515-542. <https://DOI:10.4208/aamm.2016.m-s1>
- [6] Andreianov, B. and Bouhsiss, F. Uniqueness for an Elliptic-Parabolic Problem with Neumann Boundary Condition, *Journal of Evolution Equations*, **4**, (2004), 273-295. https://doi.org/10.1007/978-3-0348-7924-8_37
- [7] A. Michel, J. Vovelle, Entropy formulation for parabolic degenerate equations with general Dirichlet boundary conditions and application to the convergence of FV methods, *Society for Industrial and Applied Mathematics Journal on Numerical Analysis*, **41**, No. 6, (2003), 2262-2293. <https://doi.org/10.1137/S0036142902406612>
- [8] R. Eymard, T. Gallouët, R. Herbin and A. Michel Convergence of a finite volume scheme for nonlinear degenerate parabolic equations, *Numerische Mathematik*, **92**, (1), (2002), 41-82. <https://doi:10.1007/s002110100342>
- [9] C. Chanais- Hillairet, Finite volumes schemes for nonlinear hyperbolic equation. Convergence towards entropy solution and error estimate, *ESAIM: Mathematical Modelling and Numerical Analysis*, **33**, (1), (1999), 129-156. <https://doi.org/10.1051/m2an:1999109>.

- [10] R.J. Diperna, Measure-valued solutions to conservation laws, *Archive for Rational Mechanics and Analysis*, **88**, (1985), 223-270. <https://doi.org/10.1007/bf00752112>
- [11] Annegret Glitzky and Jens A. Griepentrog, Discrete Sobolev Poincaré inequalities for Voronoï finite volume approximations, *Society for Industrial and Applied Mathematics Journal on Numerical Analysis*, **48**, (2010), 372-391. <https://doi.org/10.1137/09076502X>
- [12] J. Droniou, Finite volume schemes for fully non-linear elliptic equations in divergence form, *ESAIM: Mathematical Modelling and Numerical Analysis*, **40**, (6), (2006), 1069-1100. <http://eudml.org/doc/194345>
- [13] Andreianov, B. The Semigroup Approach to Conservation Laws with Discontinuous Flux. In: *Chen, GQ., Holden, H., Karlsen, K. (eds) Hyperbolic Conservation Laws and Related Analysis with Applications. Springer Proceedings in Mathematics Statistics*, vol 49. Springer, Berlin, Heidelberg (2014), pp 1-22. https://doi.org/10.1007/978-3-642-39007-4_1
- [14] Andreianov, B. and Gazibo, M.K. Explicit Formulation for the Dirichlet Problem for Parabolic-Hyperbolic Conservation Laws, *Networks and Heterogeneous Media*, **11**, No 2, (2016), 203-222. <https://doi.org/10.3934/nhm.2016.11.203>
- [15] E.Yu. Panov, On the strong pre-compactness property for entropy solutions of a degenerate elliptic equation with discontinuous flux, *Journal of Differential Equations*, **247**, (2009), 2821-2870. <https://doi.org/10.1016/j.jde.2009.08.022>
- [16] G M. Lieberman, Boundary regularity for solutions of degenerate elliptic equation, *Nonlinear Analysis: Theory, Methods and Applications*, **12**, (11), (1988), 1203-1219. [https://doi.org/10.1016/0362-546X\(88\)90053-3](https://doi.org/10.1016/0362-546X(88)90053-3)
- [17] Ph., Bénilan, H., Toure, Sur l'équation générale $u_t = a(., u, \phi(., u)_x)_x + v$ dans L^1 . I. Etude du probleme stationnaire, *In Evolution equations (Baton Rouge, LA, 1992)*, 1995, 35-62. Dekker, New York, 1995.
- [18] Ph. Bénilan and P. Wittbold. On mild and weak solutions of elliptic-parabolic problems, *Advances in Differential Equations*, **1**, (6), (1996), 1053-1073. <https://doi.org/10.57262/ade/1366895244>
- [19] R. Eymard, T. Gallouët, and R. Herbin, Finite volume methods, *Handbook of numerical analysis, Vol. VII, North Holland, Amsterdam*, (2000), 713-1020. [https://doi.org/10.1016/S1570-8659\(00\)07005-8](https://doi.org/10.1016/S1570-8659(00)07005-8).
- [20] J. Droniou, R. Eymard, and K. S. Talbot. Convergence in $C([0, T]; L^2(\Omega))$ of weak solutions to perturbed doubly degenerate parabolic equations. *Journal of Differential Equations*, **260**, (11), (2016), 7821-7860. <https://doi.org/10.1016/j.jde.2016.02.004>
- [21] B. Andreianov, M. Bendahmane, K. H. Karlsen, Discrete duality finite volume schemes for doubly nonlinear degenerate hyperbolic-parabolic equations, *Journal of Hyperbolic Differential Equations*, **7**, No 1, (2012), 1-67. <https://doi.org/10.1142/S0219891610002062>
- [22] M. Afif and B. Amaziane, Convergence of finite volume schemes for a degenerate convection-diffusion equation arising in flows in porous media, *Computer Methods in Applied Mechanics and Engineering*, **191**, (2002), 5265-5286. [https://doi.org/10.1016/S0045-7825\(02\)00458-9](https://doi.org/10.1016/S0045-7825(02)00458-9)
- [23] B. Andreianov, M. Gutnic and P. Wittbold, Convergence of finite volume approximations for a nonlinear elliptic-parabolic problem: A "continuous" approach, *Society for Industrial and Applied Mathematics Journal on Numerical Analysis*, **42**, (2004), 228-251. <https://doi.org/10.1137/S0036142901400006>
- [24] Ph. Bénilan, Crandall, M. G. and Pazy, A., Nonlinear evolution equations in Banach spaces, *Preprint book*, (1994).
- [25] F. Andreu-Vaillou, V. Casselles, J. M. Mazón, Parabolic quasilinear Equations Minimizing Linear Growth Functionals. *Progress in Mathematics*, **223**, *Birkhäuser Verlag, Basel*, (2004), xiv +340 pp. ISBN:3-7643-6619-2
- [26] Andreianov, Boris; Gazibo, Mohamed Karimou Convergence of finite volume scheme for degenerate parabolic problem with zero flux boundary condition. *Finite Volumes for Complex Applications VII- Methods and Theoretical Aspects*, Springer Proceedings in Mathematics & Statistics, **77**, Springer Cham Heidelberg New York Dordrecht London, (2014), 303-311. https://DOI:10.1007/978-3-319-05684-5_29
- [27] Alt, H.W. and Luckhaus, S. Quasilinear Elliptic-Parabolic Differential Equations. *Mathematische Zeitschrift*, **183**, No 3, (1981), 311-341. <https://doi.org/10.1007/BF0117647>

- [28] E.Yu. Panov, On the theory of generalized entropy solutions of Cauchy problem for a first-order quasilinear equation in the class of locally integrable functions, *Izvestiya Mathematics*. **66**, (6), (2002) 1171-1218 (in Russian). <https://doi.org/10.1070/IM2002v066n06ABEH000411> Authors:
- [29] Vasseur, A. Strong Traces for Solutions of Multidimensional Scalar Conservation Laws, *Archive for Rational Mechanics and Analysis*, **160**, (2001), 181-193. <https://doi.org/10.1007/s002050100157>
- [30] E.Yu. Panov, Existence of strong traces for quasi-solutions of multi-dimensional scalar conservation laws. *Journal of Hyperbolic Differential Equations*, **4**, (2009), 729-770. <https://doi.org/10.1142/S0219891607001343>

Appendix 1

We consider here a Banach space X (in application to the problem (P) , we will take $X = L^1(\Omega)$) and the multivalued operator $A : X \times X \rightarrow X$ defined by its graph. We study the general evolution problem $u' + Au \ni h, u(0) = u_0$. In our application, A is formally defined by $Au = \text{div } f(u) - \Delta \phi(u)$ with zero-flux boundary condition. In the sequel, we suppose that the operator A is m -accretive and $u_0 \in \overline{D}(A)$. We refer to [24], for definition and to [1], [6], and Appendix 2 for proof of these properties in our concrete setting which is our final purpose. In relation with the classical notion of integral solution to the abstract evolution problem introduced in [24], [25], we consider a new notion of solution called integral-process solution which depend on an additional variable $\alpha \in (0, 1)$. The purpose here is to prove that the integral-process solution of (E) coincides with mild and integral solutions. Therefore the interest of the notion of integral-process solution resides only in the fact that it may appear from some weak convergence arguments, [26], or Appendix 2 for the example we have in mind. Let us recall the notion of mild solution. In the sequel, $\|\cdot\| = \|\cdot\|_X$ being the norm in X .

Definition 7.1 A mild solution of the abstract problem $u' + Au \ni h$ on $[0, T]$ is a function $u \in C([0, T]; X)$ such that for $\sigma > 0$ there is an σ -discretization $D_A^N(t_0, \dots, t_N, h_1, \dots, h_N)$ of $u' + Au \ni h$ on $[0, T]$ which has an σ -approximate solution v satisfying

$$\|u(t) - v(t)\| \leq \sigma \text{ for } t_0 \leq t \leq t_N. \quad (73)$$

Recall that a σ -approximate solution v of $u' + Au \ni h$ on $[0, T]$ is the solution of an σ -discretization

$$D_A^N(t_0, \dots, t_N, h_1, \dots, h_N):$$

$$\frac{v_i - v_{i-1}}{t_i - t_{i-1}} + Av_i \ni h_i, \quad i = 1, 2, \dots, N \quad (74)$$

where $h \approx \sum_{i=1}^N h_i 1_{]t_{i-1}, t_i]}$ and $|t_i - t_{i-1}| \leq \sigma$. Further, v is an σ -approximate solution of the abstract initial value problem (E) if also $t_0 = 0$ and $\|v_0 - u_0\| \leq \sigma$.

Theorem 7.2 Let A be m -accretive in $L^1(\Omega)$ and $u(0) \in \overline{D}(A)$. Then the abstract initial-value problem $u' + Au \ni h$ on $(0, T]$, $u(0) = u_0$ has a unique mild solution u on $[0, T]$. Moreover u is the unique function on $C([0, T], X)$ such that for all $(\hat{u}, z) \in A$

$$\begin{aligned} & \|u(t) - \hat{u}\| - \|u(s) - \hat{u}\| \\ & \leq \int_s^t \left[u(\tau) - \hat{u}, g(\tau) - z \right] d\tau \end{aligned} \quad (75)$$

for $0 \leq s \leq t \leq T$.

Here, $[a, b] := \lim_{\lambda \downarrow 0} \frac{\|a + \lambda b\| - \|a\|}{\lambda}$ is the bracket on X , [24]. In particular if $X = L^1$ then $[a, b]_{L^1(\Omega)} = \int_{\Omega} \text{sign}(a)b \, dx + \int_{\{a=0\}} |b| \, dx$.

For the proof, we refer to [24].

A function u satisfying (75) is called integral solution. Here, we consider a more general notion of solution which is the object of this Appendix.

Definition 7.3 Let A be an accretive operator and $g \in L^1(0, T; X)$. A function $v(t, \alpha)$ is an integral-process solution of abstract problem $v' + Av \ni g$ on $[0, T]$, $v(0, \alpha) = v_0$, if v satisfy for all $(\hat{v}, z) \in A$

$$\begin{aligned} & \int_0^1 \left(\|v(t, \alpha) - \hat{v}\| - \|v(s, \alpha) - \hat{v}\| \right) d\alpha \\ & \leq \int_0^1 \int_s^t \left[v(\tau, \alpha) - \hat{v}, g(\tau) - z \right] d\tau d\alpha \end{aligned} \quad (76)$$

for $0 < s \leq t \leq T$ and the initial condition is satisfied in the sense

$$\text{ess-} \lim_{t \downarrow 0} \int_0^1 \|v(t, \alpha) - v_0\| d\alpha = 0. \quad (77)$$

Such generalization of the notion of integral solution is a purely technical hint, indeed, we show that integral-process solutions coincide with the unique integral solution in the following sense.

Theorem 7.4 Assume that A be m -accretive in X and $u_0 \in \overline{D}(A)$, u is an integral-process solution if and only if u is independent on α and for all α , $u(\cdot, \alpha)$ coincide with the unique integral and mild solution.

The result will follow directly from the proposition given below.

Proposition 7.5 *Let A be an accretive operator. If v is an integral-process solution of $v' + Av \ni g$ on $[0, T]$, $v(0, \alpha) \equiv v_0$ and u is a mild solution of $u' + Au \ni h$ on $[0, T]$, $u(0) = u_0$ then*

$$\begin{aligned} & \int_0^1 \|u(t) - v(t, \alpha)\| d\alpha \\ & \leq \int_0^t \int_0^1 \left[u(\tau) - v(\tau, \alpha), h(\tau) - g(\tau) \right] d\tau d\alpha \\ & + \int_0^1 \|u_0 - v_0\| d\alpha \end{aligned} \tag{78}$$

for a.e. $t \in [0, T]$,

Proof. Let $u_k^n, k = 1, \dots, N_n$ be a solution of the σ_n discretization $D_A(0 = t_0^n, t_1^n, \dots, t_{N_n}^n)$ of $u' + Au \ni h$ on $[0, T]$. Set $\delta_k^n = t_k^n - t_{k-1}^n$ and let $0 \leq a \leq b \leq T$. Since v is an integral-process solution of $v'(t, \alpha) + Av \ni g$ we have:

$$\begin{aligned} & \int_0^1 \left(\|v(b, \alpha) - u_k^n\| - \|v(a, \alpha) - u_k^n\| \right) d\alpha \\ & \leq \int_0^1 \int_a^b \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n + \frac{u_k^n - u_{k-1}^n}{\delta_k^n} \right] \\ & \leq \int_0^1 \int_a^b \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n \right] d\alpha d\tau \\ & + \frac{1}{\delta_k^n} \int_0^1 \int_a^b \left(\|v(\tau, \alpha) - u_{k-1}^n\| - \|v(\tau, \alpha) - u_k^n\| \right). \end{aligned} \tag{79}$$

Where we have used the inequality

$$\begin{aligned} & \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n + \frac{u_k^n - u_{k-1}^n}{\delta_k^n} \right] \\ & \leq \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n \right] \\ & + \frac{1}{\delta_k^n} \left(\|v(\tau, \alpha) - u_{k-1}^n\| - \|v(\tau, \alpha) - u_k^n\| \right) \end{aligned}$$

which follows from the facts that $\left[X, Y + Z \right] \leq \left[X, Y \right] + \left[X, Z \right]$; $\left[X, eY \right] = e \left[X, Y \right]$ if $e > 0$ and $\left[X, Y \right] \leq \frac{\|X + eY\| - \|X\|}{e}$. Multiplying (79) by δ_k^n

and summing over $k = j + 1, j + 2, \dots, i$ we find that:

$$\begin{aligned} & \sum_{k=j+1}^i \int_0^1 \delta_k^n \left(\|v(b, \alpha) - u_k^n\| - \|v(a, \alpha) - u_k^n\| \right) d\alpha \\ & \leq \sum_{k=j+1}^i \delta_k^n \int_0^1 \int_a^b \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n \right] d\tau d\alpha \\ & + \int_0^1 \int_a^b \left(\|v(\tau, \alpha) - u_j^n\| - \|v(\tau, \alpha) - u_i^n\| \right) d\tau d\alpha. \end{aligned} \tag{80}$$

Next, we assume that $\sigma_n \rightarrow 0$ and the σ_n - approximate solution of $u' + Au \ni h$ locally converge uniformly to the mild solution u on $[0, T]$. Set

$$\phi_n(t, \lambda, \alpha) = \|v(t, \alpha) - u_{k(\lambda)}^n\| \text{ for } 0 \leq t \leq T;$$

where $k(\lambda)$ is defined by $t_{k(\lambda)-1}^n < \lambda \leq t_{k(\lambda)}^n$.

Then $\phi_n(t, \lambda, \alpha) \rightarrow \|v(t, \alpha) - u(\lambda)\|$ uniformly on $[0, T] \times [0, T] \times [0, 1]$. Hence

$$\begin{aligned} & \left| \|v(t, \alpha) - u(\lambda)\| - \|v(t, \alpha) - u_{k(\lambda)}^n\| \right| \\ & \leq \|u_{k(\lambda)}^n - u(\lambda)\| \rightarrow 0. \end{aligned}$$

Therefore, if we choose i, j depending on n so that $t_j^n \rightarrow c, t_i^n \rightarrow d$ as $n \rightarrow \infty$ we have for $t \in [0, T]$

$$\sum_{k=j+1}^i \delta_k^n \|v(t, \alpha) - u_k^n\| \rightarrow \int_c^d \|v(t, \alpha) - u(\lambda)\| d\lambda.$$

Moreover with $t = \tau$, we get

$$\int_a^b \|v(\tau, \alpha) - u_j^n\| d\tau \rightarrow \int_a^b \|v(\tau, \alpha) - u(c)\| d\tau \text{ and} \tag{81}$$

$$\int_a^b \|v(\tau, \alpha) - u_i^n\| d\tau \rightarrow \int_a^b \|v(\tau, \alpha) - u(d)\| d\tau. \tag{82}$$

From now, let

$$\begin{aligned} F_n(\lambda, \alpha) &= \int_a^b \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n \right] d\tau \text{ and} \\ F(\lambda, \alpha) &= \int_{t_{k-1}^n}^{t_k^n} \int_a^b \left[v(\tau, \alpha) - u_k^n, g(\tau) - h(\lambda) \right] d\tau d\lambda \\ & \text{for } t_{k-1}^n < \lambda \leq t_k^n. \end{aligned}$$

Then

$$\begin{aligned} & \left| \sum_{k=j+1}^i \left(\delta_k^n F_n(\lambda, \alpha) - F(\lambda, \alpha) \right) \right| \\ & \leq \sum_{k=j+1}^i \int_{t_{k-1}^n}^{t_k^n} \int_a^b \|h_k^n - h(\lambda)\| d\tau d\lambda \leq \sigma_n(b - a). \end{aligned}$$

and therefore

$$\overline{\lim}_{n \rightarrow \infty} \sum_{k=j+1}^i \delta_k^n F_n(\lambda, \alpha) = \overline{\lim}_{n \rightarrow \infty} \sum_{k=j+1}^i F(\lambda, \alpha). \quad (83)$$

Since $u_k^n \rightarrow u(\lambda)$ and $t_k^n \rightarrow \lambda$ as $n \rightarrow \infty$ and the bracket $\left[\cdot, \cdot \right]$ is the upper-semicontinuous, we deduce from (83) that

$$\begin{aligned} & \overline{\lim}_{n \rightarrow \infty} \sum_{k=j+1}^i \delta_k^n \int_a^b \left[v(\tau, \alpha) - u_k^n, g(\tau) - h_k^n \right] d\tau \\ &= \overline{\lim}_{n \rightarrow \infty} \int_{t_j^n}^{t_i^n} F_n(\lambda, \alpha) d\lambda \\ &\leq \int_c^d \int_a^b \left[v(\tau, \alpha) - u(\lambda), g(\tau) - h(\lambda) \right] d\tau d\lambda. \quad (84) \end{aligned}$$

As previously, the convergence is uniform in $\alpha \in [0, 1]$, therefore we can integrate in α under the limit in (81), (82), (84) and obtain

$$\begin{aligned} & \int_0^1 \int_c^d \left(\|v(b, \alpha) - u(\lambda)\| - \|v(a, \alpha) - u(\lambda)\| \right) d\lambda d\alpha \\ &\leq \int_0^1 \int_c^d \int_a^b \left[v(\tau, \alpha) - u(\lambda), g(\tau) - h(\lambda) \right] d\tau d\lambda d\alpha \\ &+ \int_0^1 \int_c^d \left(\|v(\tau, \alpha) - u(c)\| - \|v(\tau, \alpha) - u(d)\| \right) d\tau d\alpha. \quad (85) \end{aligned}$$

Now, we set:

$$\begin{aligned} \varpi(s, t, \alpha) &= \int_0^1 \|v(s, \alpha) - u(t)\| d\alpha \\ \Pi(s, t, \alpha) &= \int_0^1 \left[v(s, \alpha) - u(t), g(s) - h(t) \right] d\alpha. \end{aligned}$$

Recall that $u \in C([0, T]; X)$ and u_0 satisfies $\text{ess-}\lim_{t \downarrow 0} \int_0^1 \|v(t, \alpha) - u_0\| d\alpha = 0$. Then, v is continuous a.e. for any Lebesgue point on $[0, T]$. The function ϖ and Π are continuous in t and integrable in s

$$\varphi(t, t) - \varphi(s, s) \leq \int_0^t \Pi(\tau) d\tau - \int_0^s \Pi(\tau) d\tau.$$

Then for a.e. $t, s \in [0, T]$

$$\begin{aligned} \Xi(t) &= \varphi(t, t) - \int_0^t \Pi(\tau) d\tau \\ &\leq \varphi(s, s) - \int_0^s \Pi(\tau) d\tau = \Xi(s) \text{ for a.e. } t, s \in [0, T]. \end{aligned}$$

The function Ξ is continuous at 0^+ , therefore $\Xi(t) \leq \Xi(0)$. This is equivalent to (78).

Appendix 2

In this appendix, we apply the notion of integral-process solution to the problem (P) and present a way to prove uniqueness of entropy solution. In [7], the authors introduced a notion of entropy-process solution and using the doubling of variables method of [3], they proved that entropy solution is the unique entropy-process solution. In our case, we were not able to use the same argument because we need that the entropy solution possess a strong boundary trace on the boundary in order that the doubling of variables apply for example refer to [6]. Fortunately, under additional assumptions, we can ensure the desired boundary regularity for the associated stationary problem:

$$(S) \begin{cases} v + \text{div}(f(v) - \nabla\phi(v)) = g & \text{in } \Omega, \\ (f(v) - \nabla\phi(v)) \cdot \eta = 0 & \text{on } \partial\Omega. \end{cases}$$

Therefore, firstly we compare an entropy-process solution μ of (P) with the entropy solution of (S). This suggests the use of nonlinear semigroup theory; more precisely we find that μ is also an integral-process solution to $u' + Au = 0, \mu(0, \alpha) = u_0$ with appropriately defined operator A . Then, proving the m-accretivity of A and using the Appendix 1 we are able to conclude that μ is the unique mild and integral solution of the abstract evolution problem. At the last step, we use the result of [1], which says that such solution is the unique entropy solution of (P).

Proposition 8.1 *Let $\xi \in C^\infty([0, T] \times \mathbb{R}^\ell), \xi \geq 0$. Then for all $k \in]u_c, u_{\max}]$, for all $D \in \mathbb{R}^\ell$ and for all entropy-process solution μ of (P), we have*

$$\begin{aligned} & \int_0^T \int_\Omega \int_0^1 |\mu - k| \xi_t dx dt d\alpha + \int_\Omega |u_0 - k| \xi(0, x) dx \\ &+ \int_0^T \int_\Omega \int_0^1 \text{sign}(\mu - k) (f(\mu) - f(k)) \cdot \nabla \xi dx dt d\alpha \\ &- \int_0^T \int_\Omega \text{sign}(u - k) \left(\nabla \phi(u) - D \right) \cdot \nabla \xi dx dt \\ &+ \int_0^T \int_{\partial\Omega} |(f(k) - D) \cdot \eta(x)| \xi d\mathcal{H}^{\ell-1}(x) dt \\ &\geq \overline{\lim}_{\sigma \rightarrow 0} \frac{1}{\sigma} \int \int_{Q \cap A_k^\sigma} \nabla \phi(u) \cdot \left(\nabla \phi(u) - D \right) \xi dx dt. \quad (86) \end{aligned}$$

with $A_k^\sigma = \{-\sigma < \phi(u) - \phi(k) < \sigma\}$.

Proof. The proof follows the arguments of [1]. Let us recall that if $\mu(t, x, \alpha)$ is entropy-process solution and (1) holds, then $u(t, x) = \int_0^1 \mu(t, x, \alpha) d\alpha$ satisfies in

the weak sense for all $k \in [u_c, u_{\max}]$ and all $D \in \mathbb{R}^\ell$:

$$(Pk) \begin{cases} (u - k)_t + \operatorname{div}(\mathcal{F}[\mu] - \mathcal{F}[k]) = 0 \text{ in } Q, \\ \operatorname{ess-} \lim_{t \rightarrow 0^+} (u(t, x) - k) = u_0(x) - k \text{ on } \Omega, \\ (\mathcal{F}[\mu] - \mathcal{F}[k]) \cdot \eta = -\mathcal{F}[k] \cdot \eta \text{ on } \Sigma. \end{cases}$$

where

$$\mathcal{F}[u] = \int_0^1 f(\mu) d\alpha - \nabla \phi(u)$$

$$\mathcal{F}[k] = f(k) - D.$$

Take the test function $\operatorname{sign}_\sigma(\phi(u) - \phi(k))\xi = H_\sigma(\phi(u) - \phi(k))\xi$ in the weak formulation of this problem with $\xi \in C^\infty([0, T] \times \mathbb{R}^\ell)$. Using the formalism of [27], we have

$$\begin{aligned} & \int_0^T \langle (u - k)_t, H_\sigma(\phi(u) - \phi(k))\xi \rangle_{H^1(\Omega)^*, H^1(\Omega)} dt \\ & - \int_0^T \int_\Omega H_\sigma(\phi(u) - \phi(k)) (\mathcal{F}[\mu] - \mathcal{F}[k]) \cdot \nabla \xi \\ & - \int_0^T \int_\Omega \xi (\mathcal{F}[\mu] - \mathcal{F}[k]) \cdot \nabla H_\sigma(\phi(u) - \phi(k)) \\ & - \int_0^T \int_{\partial\Omega} H_\sigma(\phi(u) - \phi(k)) \mathcal{F}[k] \cdot \eta \xi = 0. \end{aligned} \quad (87)$$

By the weak chain rule in the sense of [27]

$$\begin{aligned} & \int_0^T \langle (u - k)_t, H_\sigma(\phi(u) - \phi(k))\xi \rangle_{H^1(\Omega)^*, H^1(\Omega)} \\ & = - \int_0^T \int_\Omega I_\sigma(u)\xi_t dt dx - \int_\Omega I_\sigma(u_0)\xi(0, x) dx \end{aligned}$$

where: $H_\sigma(r) = \begin{cases} 1 & \text{if } r > \sigma, \\ \frac{r}{\sigma} & \text{if } |r| \leq \sigma, \\ -1 & \text{if } r < -\sigma, \end{cases}$ and

$$I_\sigma : z \mapsto \int_k^z H_\sigma(\phi(s) - \phi(k)) ds \rightarrow |z - k|$$

as $\sigma \rightarrow 0$.

Then, after passing to the limit as $\sigma \rightarrow 0$, we have

$$\begin{aligned} & \int_Q |u - k| \xi_t dx dt \\ & = - \int_Q \operatorname{sign}(u - k)(u - k) \xi_t dx dt \\ & = - \int_Q \operatorname{sign}(u - k) \left(\int_0^1 \mu d\alpha - k \right) \xi_t dx dt \end{aligned}$$

Now, notice that because $k \in]u_c, u_{\max}[$ and because $\phi(\mu(\alpha)) = \operatorname{const}$ on $[0, 1]$ we find that $\operatorname{sign}(\mu(\alpha) - k)$ is constant on $[0, 1]$ equal to $\operatorname{sign}(u - k)$. Then, we see that

$$\int_Q |u - k| \xi_t dx dt = - \int_Q \int_0^1 |\mu - k| \xi_t d\alpha dx dt$$

Similarly, we see

$$\begin{aligned} & \int_0^T \int_\Omega \int_0^1 \operatorname{sign}(u - k) [f(\mu) - f(k)] \cdot \nabla \xi dx dt \\ & = \int_0^T \int_\Omega \int_0^1 \operatorname{sign}(\mu - k) [f(\mu) - f(k)] \cdot \nabla \xi dx dt d\alpha \end{aligned}$$

For treatment of the others terms, we refer to [1].

Let us firstly prove that the initial datum is satisfied in the sense of (77) (Appendix 1). This means that the entropy-process solution satisfies the initial condition of integral-process solution.

Lemma 8.2 *Let v be an entropy-process solution of (P) with initial datum $v_0 \in L^\infty$. Then the initial datum is taken in the following sense:*

$$\lim_{s \downarrow 0} \int_0^s \int_\Omega \int_0^1 |v - v_0| dt dx d\alpha = 0. \quad (88)$$

Proof. The proof follows the idea of Panov in ([28], Proposition 1). For $c \in \mathbb{R}$ and $s > 0$, consider the functions

$$\Psi_s(\cdot, c) : x \in \Omega \mapsto \frac{1}{s} \int_0^s \int_0^1 |v(t, x, \alpha) - c| dt d\alpha. \quad (89)$$

Because v is bounded, the set $(\Psi_s(\cdot, c))_{s>0}$ is bounded in $L^\infty(\Omega)$. Therefore for any sequence $s_m \rightarrow 0$, there exists a subsequence such that for all $c \in \mathbb{Q}$, $(\Psi_{s_m}(\cdot, c))_{s_m>0}$ converges in $L^\infty(\Omega)$ weak star to some limit denoted by $(\Psi(\cdot, c))$. Fix $\xi \in \mathcal{D}(\Omega)^+$. From Remark 6.3 with test function $\tilde{\xi}(t, x) := (1 - \frac{t}{s})^+ \xi(x)$ we readily infer the inequalities $\forall c \in \mathbb{Q}$:

$$\int_\Omega \Psi(x, c) \xi(x) dx \leq \int_\Omega |u_0 - c| \xi(x) dx. \quad (90)$$

By the density argument, we extend (90) to all $\xi \in L^1(\Omega)$, $\xi \geq 0$. Now for all $\epsilon > 0$, there exists a number $N(\epsilon) \in \mathbb{N}$, a collection $(c_j^\epsilon)_{j=1}^{N(\epsilon)} \subset \mathbb{Q}$ and a partition of Ω into disjoint union of measurable sets $\Omega_1^\epsilon, \dots, \Omega_{N(\epsilon)}^\epsilon$ such that $\|v_0 - v_0^\epsilon\|_{L^1} \leq \epsilon$, where $v_0^\epsilon := \sum_{j=1}^{N(\epsilon)} c_j^\epsilon 1_{\Omega_j^\epsilon}$. Because $1_\Omega = \sum_{j=1}^{N(\epsilon)} 1_{\Omega_j^\epsilon}$, apply-

ing (90) with $c = c_j^\epsilon$ and $\xi = 1_{\Omega_j^\epsilon}$ we deduce

$$\begin{aligned} & \lim_{m \rightarrow \infty} \frac{1}{s_m} \int_0^{s_m} \int_{\Omega} \int_0^1 |v - v_0^\epsilon| dt dx d\alpha \\ &= \lim_{m \rightarrow \infty} \int_{\Omega} \sum_{j=1}^{N(\epsilon)} \Psi_{s_m}(x, c_j^\epsilon) 1_{\Omega_j^\epsilon} dx \\ &= \int_{\Omega} \sum_{j=1}^{N(\epsilon)} \Psi(x, c_j^\epsilon) 1_{\Omega_j^\epsilon} dx \\ &\leq \int_{\Omega} \sum_{j=1}^{N(\epsilon)} |v_0 - c_j^\epsilon| 1_{\Omega_j^\epsilon} dx = \|v_0 - v_0^\epsilon\|_{L^1} \leq \epsilon. \end{aligned}$$

Using once more the bound $\|v_0 - v_0^\epsilon\|_{L^1} \leq \epsilon$ (in the first term of the previous calculation), we can send ϵ to zero and infer the analogue of (89), with a limit taken along some subsequence of $(s_m)_{m>1}$. Because $(s_m)_{m>1}$ was an arbitrary sequence convergent to zero, (89) is justified.

Now it remains to prove that the entropy-process solution an is integral-process solution. Let us define the (possibly multivalued) operator L_f^ϕ by it resolvent

$$(v, z) \in L_f^\phi = \left\{ \begin{array}{l} v \text{ such that } v \text{ is an entropy solution} \\ \text{of } (S), \text{ with } g = v + z. \\ \text{and strong } L^1 \text{ trace of} \\ (f(u) - \nabla\phi(u)) \cdot \eta|_{\partial\Omega} \text{ exists} \\ \text{and equal to zero} \end{array} \right\}$$

Definition 8.3 *The normal component of the flux $\mathcal{F}[u] = (f(u) - \nabla\phi(u)) \cdot \eta$ has a L^1 strong trace $\gamma\mathcal{F}[\hat{u}] \in L^1_{Loc}(\partial\Omega)$, at boundary $\partial\Omega$ if*

$$\lim_{s \rightarrow 0} \frac{1}{s} \int_0^s \int_{\hat{x} \in \partial\Omega} \xi(\hat{x}) |\mathcal{F}[u](s, \hat{x}) - \gamma\mathcal{F}[u](\hat{x})| d\hat{x} d\tau = 0.$$

After having defined the operator L_f^ϕ , we present the following results.

Theorem 8.4 *Assume that L_f^ϕ is m -accretive densely defined on $L^1(\Omega; [0, u_{\max}])$. Then the entropy-process solution is the unique entropy solution.*

Before turning to the proof of Theorem 8.4, lets us present three cases where it applies.

Proposition 8.5 *Assume (66) holds. Then, L_f^ϕ is m -accretive densely defined on $L^1(\Omega; [0, u_{\max}])$.*

For the proof, we refer to [1], Proposition 4.10.

Proposition 8.6 *Assume that, (67), holds. Then L_f^ϕ is m -accretive densely defined on $L^1(\Omega; [0, u_{\max}])$.*

Proof.(sketch) The proof is essentially the same as in [6], where the case $\phi = Id$ has been investigated. For general ϕ satisfying $f \circ \phi^{-1} \in C^{0,\alpha}$, $\alpha > 0$ we adapt the result of [16]. As ϕ is bijective, we set $w = \phi(u)$ and rewrite the stationary problem as:

$$\begin{aligned} \operatorname{div}(f \circ \phi^{-1}(w) - \nabla w) &= g(x) - \phi^{-1}(w) \\ \Rightarrow \operatorname{div}(B(w, \nabla w)) &= F(x, w), \end{aligned}$$

where B and F satisfies the hypothesis of [16], then $w = \phi(u) \in C^{0,\alpha}(\bar{\Omega})$, $\alpha > 0$ and $u \in C^{0,\alpha}(\bar{\Omega})$. We deduce that $(f(u) - \nabla(\phi(u))) \in C(\bar{\Omega})$.

Proposition 8.7 *Assume that (68) holds. Then L_f^ϕ is m -accretive densely defined on $L^1(\Omega; [0, u_{\max}])$.*

For the proof, we refer to [4], [29], [30], where the existence of strong trace of $f(u)$ has been proved for pure conservation laws. In the sequel, we concentrate on the proof of Theorem 8.4 in the case (66) holds. The other cases are similar, using the hint of [6].

Proof of Theorem 8.4 Now, we apply the doubling of variables, [3], in the way of [6], [1]. We consider $\mu = \mu(t, x, \alpha)$ an entropy-process solution of (P) and $v = v(y)$ an entropy solution of (S) using in the definition of L_f^ϕ . Consider non-negative function $\xi = \xi(t, x, y)$ having the property that $\xi(\cdot, \cdot, y) \in C^\infty([0, T] \times \bar{\Omega})$ for each $y \in \bar{\Omega}$, $\xi(t, x, \cdot) \in C_0^\infty(\bar{\Omega})$ for each $(t, x) \in [0, T] \times \bar{\Omega}$. Let us denote the sets on which the diffusion term for the first, respectively for the second solutions degenerate by $\Omega_x = \{x \in \Omega; \mu(t, x, \alpha) \in [0, u_c]\}$; $\Omega_y = \{y \in \Omega; v(y) \in [0, u_c]\}$. We denote by Ω_x^c respectively Ω_y^c their complementaries in Ω . In (86), take $\xi = \xi(t, x, y)$, $k = u(y)$, $D = \phi(u)_y$ and integrate

over $\Omega_y^c \times [0, 1]$. We get

$$\begin{aligned} & \int_{\Omega_y^c} \int_0^T \int_{x \in \Omega} \int_0^1 |\mu - v| \xi_t d\alpha dx dt dy \\ & + \int_{\Omega_y^c} \int_0^T \int_{x \in \Omega} \int_0^1 \text{sign}(\mu - v) [f(\mu) - f(v)] \cdot \xi_x \\ & - \int_{\Omega_y^c} \int_0^T \int_{x \in \Omega} \text{sign}(u - v) (\phi(u)_x - \phi(v)_y) \cdot \xi_x \\ & + \int_{\Omega_y^c} \int_0^T \int_{x \in \partial\Omega} |(f(v) - \phi(v)_y) \cdot \eta(x)| \xi d\sigma dt dy \\ & + \int_{\Omega_y^c} \int_{x \in \Omega} |u_0 - v| \xi(0, x, y) dx dy \\ & \geq \lim_{\sigma \rightarrow 0} \frac{1}{\sigma} \int_{\Omega_y^c} \int_0^T \int_{x \in \Omega \cap A_u^\sigma} \phi(u)_x (\phi(u)_x - \phi(v)_y) \xi. \end{aligned} \tag{91}$$

In the same way, in (6.3) take $\xi = \xi(t, x, y)$, $k = v(y)$, integrate over Ω_y , and use the fact that $\phi(v)_y = 0$ in Ω_y . We get

$$\begin{aligned} & \int_{\Omega_y} \int_0^T \int_{x \in \Omega} \int_0^1 |\mu - v| \xi_t d\alpha dx dt dy \\ & + \int_{\Omega_y} \int_0^T \int_{x \in \Omega} \int_0^1 \text{sign}(\mu - v) [f(\mu) - f(v)] \cdot \xi_x \\ & - \int_{\Omega_y} \int_0^T \int_{x \in \Omega} \text{sign}(u - v) (\phi(u)_x - \phi(v)_y) \cdot \xi_x \\ & + \int_{\Omega_y} \int_0^T \int_{x \in \partial\Omega} |(f(v) - \phi(v)_y) \cdot \eta(x)| \xi d\sigma dt dy \\ & + \int_{\Omega_y} \int_{x \in \Omega} |u_0 - v| \xi(0, x, y) dx dy \geq 0. \end{aligned} \tag{92}$$

Since $\Omega = \Omega_x \cup \Omega_x^c$, by adding (91) to (92) we obtain:

$$\begin{aligned} & \int_{y \in \Omega} \int_0^T \int_{x \in \Omega} \int_0^1 |\mu - v| \xi_t d\alpha dx dt dy \\ & + \int_{y \in \Omega} \int_0^T \int_{x \in \Omega} \text{sign}(\mu - v) [f(\mu) - f(v)] \cdot \xi_x \\ & - \int_{y \in \Omega} \int_0^T \int_{x \in \Omega} \text{sign}(u - v) (\phi(u)_x - \phi(v)_y) \cdot \xi_x \\ & + \int_{\Omega} \int_0^T \int_{x \in \partial\Omega} |(f(v) - \phi(v)_y) \cdot \eta(x)| \xi d\sigma dt dy \\ & + \int_{\Omega} \int_{x \in \Omega} |u_0 - v| \xi(0, x, y) dx dy \\ & \geq \lim_{\sigma \rightarrow 0} \frac{1}{\sigma} \int_{\Omega_y^c} \int_0^T \int_{x \in \Omega \cap A_u^\sigma} \phi(u)_x (\phi(u)_x - \phi(v)_y) \xi. \end{aligned} \tag{93}$$

In the entropy formulation of (S), take $\xi = \xi(t, x, y)$, $k = \mu(t, x, \alpha)$, $D = \phi(\mu)_x$ and integrate over $(t, x, \alpha) \in (0, T) \times \Omega_x^c \times (0, 1)$

$$\begin{aligned} & \int_{\Omega_x^c} \int_0^T \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) [f(v) - f(\mu)] \cdot \xi_y \\ & - \int_{\Omega_x^c} \int_0^T \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) (\phi(v)_y - \phi(u)_x) \cdot \xi_y \\ & + \int_0^T \int_{\Omega_x^c} \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) (v - g(y)) \xi \\ & + \int_0^T \int_{\Omega_x^c} \int_0^1 \int_{y \in \partial\Omega} |(f(\mu) - \phi(\mu)_x) \cdot \eta(y)| \xi \\ & \geq \lim_{\sigma \rightarrow 0} \frac{1}{\sigma} \int_{\Omega_x^c} \int_0^T \int_{y \in \Omega \cap A_v^\sigma} \int_0^1 \phi(v)_y (\phi(v)_y - \phi(\mu)_x) \xi. \end{aligned} \tag{94}$$

Since $v(y)$ is entropy solution, then take in the entropy dissipative formulation of (S) $\xi = \xi(t, x, y)$, $k = \mu(t, x, \alpha) \in]u_c, u_{\max}[$, integrate over $(0, T) \times \Omega_x \times (0, 1)$ and use the fact that $\phi(\mu)_x = \phi(u)_x = 0$ in $(0, T) \times \Omega_x$.

$$\begin{aligned} & \int_{\Omega_x} \int_0^T \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) [f(v) - f(\mu)] \cdot \xi_y \\ & - \int_{\Omega_x} \int_0^T \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) (\phi(v)_y - \phi(u)_x) \cdot \xi_y \\ & + \int_0^T \int_{\Omega_x} \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) (v - g(y)) \xi \\ & + \int_0^T \int_{\Omega_x} \int_0^1 \int_{y \in \partial\Omega} |(f(\mu) - \phi(\mu)_x) \cdot \eta(y)| \xi \geq 0 \end{aligned} \tag{95}$$

By adding (94) to (95), we obtain

$$\begin{aligned} & \int_{\Omega} \int_0^T \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) [f(v) - f(\mu)] \cdot \xi_y \\ & - \int_{\Omega} \int_0^T \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) (\phi(v)_y - \phi(u)_x) \cdot \xi_y \\ & + \int_0^T \int_{\Omega} \int_0^1 \int_{y \in \Omega} \text{sign}(v - \mu) (v - g(y)) \xi \\ & + \int_0^T \int_{\Omega} \int_0^1 \int_{y \in \partial\Omega} |(f(\mu) - \phi(\mu)_x) \cdot \eta(y)| \xi \\ & \geq \lim_{\sigma \rightarrow 0} \frac{1}{\sigma} \int_{\Omega_x^c} \int_0^T \int_{\Omega_y^c \cap A_u^\sigma} \int_0^1 \phi(v)_y (\phi(v)_y - \phi(\mu)_x) \xi. \end{aligned} \tag{96}$$

Now, sum (93) and (96) to obtain

$$\begin{aligned}
 & \int_0^1 \int_0^T \int_{\Omega} \int_{\Omega} |\mu - v| \xi_t d\alpha dy dx dt \\
 & + \int_{\Omega} \int_{\Omega} |u_0 - v| \xi(0, x, y) dx dy \\
 & + \int_0^1 \int_0^T \int_{\Omega} \int_{\Omega} \text{sign}(v - u) [f(\mu) - f(v)] \cdot \xi_{x+y} \\
 & - \int_0^T \int_0^1 \int_{\Omega} \int_{\Omega} \text{sign}(u - v) \left(\phi(u)_x - \phi(v)_y \right) \cdot \xi_{x+y} \\
 & + \int_0^T \int_{x \in \partial\Omega} \int_{\Omega} |(f(v) - \phi(v)_y) \cdot \eta(x)| \xi d\sigma dt dy \\
 & + \int_0^1 \int_0^T \int_{\Omega} \int_{y \in \partial\Omega} |(f(\mu) - \phi(\mu)_x) \cdot \eta(y)| \xi dy d\sigma dt \\
 & + \int_0^1 \int_0^T \int_{\Omega} \int_{\Omega} \text{sign}(v - \mu) (v - g(y)) \xi dy dx dt d\alpha \\
 & \geq \overline{\lim}_{\sigma \rightarrow 0} \frac{1}{\sigma} \int_0^T \int_{\Omega_x^c \times \Omega_y^c \cap A_{\mu}^{\sigma}} |\phi(v)_x - \phi(u)_y|^2 \xi dy dx dt \\
 & \geq 0.
 \end{aligned} \tag{97}$$

Next, following the idea of [6], in the simple one-dimensional setting, we consider the test function $\xi(t, x, y) = \theta(t) \rho_n(x, y)$, where $\theta \in C_0^{\infty}(0, T)$, $\theta \geq 0$, $\rho_n(x, y) = \delta_n(\Delta)$ and $\Delta = (1 - \frac{1}{n(b-a)})x - y + \frac{a+b}{2n(b-a)}$. Then, $\rho_n \in \mathcal{D}(\overline{\Omega} \times \overline{\Omega})$ and $\rho_n|_{\Omega \times \partial\Omega} = 0$. Due to this choice

$$\int_0^T \int_{x \in \Omega} \int_{y \in \partial\Omega} \int_0^1 |(f(\mu) - \phi(u)_x) \cdot \eta(y)| \rho_n \theta = 0.$$

By the Proposition 8.5 and the definition of L_f^{ϕ} , we prove that for the stationary problem, $(f(v) - \phi(v)_y) \in C_0([a, b])$. Therefore we have $|(f(v) - \phi(v)_y) \cdot \eta(x)| \rightarrow 0$ when $x \rightarrow y$, i.e, as $n \rightarrow \infty$. We conclude that

$$\lim_{n \rightarrow \infty} \int_0^T \int_{x \in \partial\Omega} \int_{y \in \Omega} |(f(v) - \phi(v)_y) \cdot \eta(x)| \rho_n \theta = 0.$$

It remains to study the limit, as $n \rightarrow \infty$

$$I_n = \int_0^1 \int_0^T \int_{\Omega} \int_{\Omega} \theta \text{sign}(\mu - v) F(\mu, v) \cdot \left((\rho_n)_x + (\rho_n)_y \right).$$

we proved in [1], that the limit of I_n equals zero. The passage to the limit in other terms in (97) is straightforward. Finally (97) gives for $n \rightarrow \infty$

$$\begin{aligned}
 & \int_0^1 \int_0^T \int_{\Omega} |\mu(t, x, \alpha) - v(y)| \theta'(t) dx dt d\alpha \\
 & + \int_0^1 \int_0^T \int_{\Omega} \text{sign}(v - \mu) (v - g) \theta dx dt d\alpha \geq 0.
 \end{aligned}$$

Hence

$$\begin{aligned}
 & \frac{d}{dt} \int_0^1 \|\mu(t, \alpha) - v\|_{L^1(\Omega)} d\alpha \\
 & \leq \int_0^1 \int_{\Omega} \text{sign}(\mu - v) (v - g) dx \text{ in } \mathcal{D}'(0, T).
 \end{aligned}$$

Thus, μ is an integral-process solution of (E) with $A = L_f^{\phi}$. Now, the claim of Theorem 8.4 is a direct consequence of the fact that the integral-process solution is a unique integral solution (Appendix 1) and then is an entropy solution of (P).

Contribution of Individual Authors to the Creation of a Scientific Article (Ghostwriting Policy)

The authors equally contributed in the present research, at all stages from the formulation of the problem to the final findings and solution.

Sources of Funding for Research Presented in a Scientific Article or Scientific Article Itself

No funding was received for conducting this study.

Conflict of Interest

The authors have no conflicts of interest to declare that are relevant to the content of this article.

Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

https://creativecommons.org/licenses/by/4.0/deed.en_US