

- [4] T. N. Land and N. T. Binh, "Daily roadside BTEX concentrations in East Asia measured by the Lanwatsu, Radiello and Ultra I SKS passive samplers," *Science of the Total Environment*, vol. 441, pp. 248-257, 2012.
- [5] J. G. Cerón Bretón, R. M. Cerón Bretón, J. D. W. Kahl, E. Ramírez Lara, C. Guarnaccia, C. A. Aguilar Uacán and U. López Chuken, "Diurnal and seasonal variation of BTEX in the air of Monterrey, Mexico: preliminary study of sources and photochemical ozone pollution.," *Air Quality, Atmosphere & Health*, vol. 8(5), pp. 469-482, 2014.
- [6] J. G. Cerón Bretón, R. M. Cerón Bretón, F. V. Ucan, C. B. Baeza, M. D. L. L. E. Fuentes, E. R. Lara, and M. U. Chi, "Characterization and sources of Aromatic Hydrocarbons (BTEX) in the atmosphere of two urban sites located in Yucatan Peninsula in Mexico," *Atmosphere*, vol. 107, p. 8(6), 2017.
- [7] Z. Zhang, X. Wang, Y. Zhang, S. Lu, X. Huang and Y. Wang, "Ambient air benzene at background sites in China's most developed coastal regions: Exposure levels, source implications and health risks.," *Science of the Total Environment*, vol. 511, pp. 792-800, 2015.
- [8] Ceron, J.G.; Ramirez, E.; Ceron, R.M.; Carballo, C.; Aguilar, C.; Lopez, U.; Ramirez, A.; Garcias, Y.; Naal, D.; Campero, A.; et al., "Diurnal and Seasonal Variation of BTX in Ambient air of One Urban Site in Carmen City, Campeche, Mexico", *J. Environ. Prot.* Vol. 4, pp. 40–49, 2013.
- [9] Ceron-Breton, J.G., Ceron-Breton, R.M., Martinez-Morales, S., Kahl, J.D.W., Guarnaccia, C., Lara-Severino, R.C., Rangel-Marron, M., Ramirez-Lara, E., Espinosa-Fuentes, M.L., Uc-Chi, M.P., and Licona-Sanchez, G, "Health Risk Assessment of the levels of BTEX in ambient air of one urban site located in Leon, Guanajuato, Mexico during two climatic seasons", *Atmosphere*, vol. 11 (165), pp. 1-22, 2020.
- [10] Bravo, A.H., E.R. Sosa, A.P. Sánchez, L.E. Bueno, & R.L. González, "Concentrations of benzene and toluene in the atmosphere of the southwestern area at the Mexico City Metropolitan Zone", *Atmos. Environ.*, vol. 36, pp. 3843-3849, 2002.
- [11] Zhang, Y.; Mu, Y.; Liu, J.; Mellouki, A., "Levels, sources and health risks of carbonyls and BTEX in the ambient air of Beijing, China", *J. Environ. Sci.* vol. 24, pp. 124–130, 2012.
- [12] U.S. Environmental Protection Agency (EPA), "Integrated Risk Information System (IRIS). Toxicological review of benzene (non-cancer effects) (EPA/635/R-02/001F)," 2014. [Online]. Available: <http://www.epa.gov/iris/tox-reviews/0276tr.pdf>.
- [13] B. Pekey and H. Yılmaz, "The use of passive sampling to monitor spatial trends of volatile organic compounds (VOCs) at an industrial city of Turkey.," *Microchemical Journal*, vol. 97(2), p. 213–219, doi: 10.1016/j.microc.2010.09.006, 2011.
- [14] D. Estéves, N. E. Dimas, G. M. T. Cassanova, J. G. Cerón Bretón, R. M. Cerón Bretón, P. E. Rustián, E. Houbron and C. Cima, "Niveles de BTEX en aire ambiente y niveles de riesgo carcinogénico de benceno en un sitio urbano del centro de Veracruz," in *2º congreso Nacional AMICA 2015*, Puebla, Puebla, 2015.
- [15] S. E. Carranco Lozada, J. G. Cerón Bretón, R. M. Cerón Bretón, A. Rodríguez Guzmán y A. R. Tovar Gerardo, «Calidad del aire en Tijuana, Baja California durante invierno de 2017,» de *CIM, Orizaba*, 2018.

Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0 https://creativecommons.org/licenses/by/4.0/deed.en_US