

Extended And Unified Local Convergence For Newton-Kantorovich Method Under w - Conditions With Applications

IOANNIS K. ARGYROS

Cameron University
Department of Mathematical Sciences
Lawton, OK 73505
USA
iargyros@cameron.edu

SANTHOSH GEORGE

NIT Karnataka
Department of Mathematical
and Computational Sciences
Surathkal
INDIA
sgeorge@nitk.ac.in

Abstract: The goal of this paper is to present a local convergence analysis of Newton's method for approximating a locally unique solution of an equation in a Banach space setting. Using the gauge function theory and our new idea of restricted convergence regions we present an extended and unified convergence theory.

Key-Words: Newton's method, Banach space, semilocal convergence, gauge function, convergence region, Newton-Kantorovich theorem.

1 Introduction

Many problems in computational sciences and related areas can be formulated as equations defined on some spaces using Mathematical modeling [2,4,5,7,12,22]. The closed form solution of such equations is desirable but rarely attainable. That is why, in practice we utilize iterative methods. There is a plethora of local as well as semi-local convergence results for Newton's method under generalized Lipschitz-type conditions [1–49]. Newton's method defined for each $n = 0, 1, 2, \dots$ by

$$x_{n+1} = N(x_n), \quad (1.1)$$

is undoubtedly the most popular method for constructing a sequence $\{x_n\}$ approximating a solution p of nonlinear equation

$$F(x) = 0. \quad (1.2)$$

Here x_0 is an initial point, $F : D \subseteq \mathcal{B}_1 \rightarrow \mathcal{B}_2$, D is a convex subset of \mathcal{B}_1 ; $\mathcal{B}_1, \mathcal{B}_2$ are Banach spaces, F is a Fréchet-differentiable operator and $N(x) = x - F'(x)^{-1}F(x)$.

Proinov in [31] presented a generalization for the famous Banach contraction mapping principle with arbitrary order of convergence of Picard's iteration. Then, later in [33] some general local convergence results were reported for Picard's iteration with arbitrary order of convergence using an iteration function in a metric space. More recently, the semi-local convergence was given in [32] with order of convergence $\gamma \geq 1$ for Picard's iteration defined for each

$n = 0, 1, 2, \dots$ by

$$x_{n+1} = P(x_n), \quad (1.3)$$

where $P : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_1$ is an iteration function in \mathcal{B}_1 satisfying the condition

$$H(P(x)) \leq g(H(x)) \text{ for each } x \in D \quad (1.4)$$

with $P(x) \in D$ and $H(x) \in I$, $H : D \rightarrow \mathbb{R}_+$, $I \subseteq \mathbb{R}_+$ containing 0 and g is a gauge function of I (to be precised in Section 2). $H : D \rightarrow \mathbb{R}_+$ satisfying (1.4) is called a function of initial approximation of P . Using g, H and P the local convergence of Picard's method was given in [31]. These results were then specialized using w -versions of the Newton-Kantorovich theorem [22]. These results extended and unified earlier ones.

In the present study, we are motivated by work in [33] and optimization considerations. Using our new idea of restricted convergence regions we find a more precise location, where the Newton iterates lie than before. This way the resulting w - functions are smaller than before leading to the following advantages (\mathcal{A}):

- (\mathcal{A}_1) Extended region of convergence leading to a wider choice of initial guesses.
- (\mathcal{A}_2) At least as precise error bounds on the distances $\|x_n - p\|$ leading to fewer iterates to obtain a desired error tolerance.
- (\mathcal{A}_3) An at least as precise information on the location of the solution p .

Advantages (\mathcal{A}) are obtained under the same computational effort, since in practice the computation of the w -function in [33] requires the computation of our new w -functions as special cases.

The rest of the study is structured as follows: Some Mathematical background on gauge functions and related materials is presented in Section 2. The local convergence of Newton’s method is given in Section 3.

2 Gauge functions and initial conditions

In order to make the study as self contained as possible, we reproduce some gauge function related concepts. More details can be found in [32, 33].

Definition 2.1 A function $g : I \rightarrow \mathbb{R}_+$ is called quasi-homogeneous of degree $\delta \geq 0$ on I if it satisfies the condition

$$g(\lambda t) \leq \lambda^\delta g(t) \text{ for each } \lambda \in (0, 1) \text{ and } t \in I. \quad (2.1)$$

Definition 2.2 A function $g : I \rightarrow \mathbb{R}_+$ is called gauge function of order $\delta \geq 1$ on I if it is quasi-homogeneous of order δ on I and

$$g(t) \leq t \text{ for each } t \in I. \quad (2.2)$$

Definition 2.3 Let $P : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$. A function $H : D \rightarrow \mathbb{R}_+$ is called function of initial conditions of P with a gauge function g on I if there exists a function $g : I \rightarrow I$ such that

$$H(P(x)) \leq g(H(x)) \text{ for each } x \in D \text{ with } P(x) \subset D \text{ and } H(x) \in I \quad (2.3)$$

Definition 2.4 Let $P : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_1$ and let $H : D \rightarrow \mathbb{R}_+$ be a function of initial conditions of P with a gauge function on I . Then a point $x \in D$ is called initial point of P if $H(x) \in I$ and iterates $P^n x$ are well-defined and remain in D .

We shall need the following auxiliary result (see corollary 3.7 in [33]).

COROLLARY 2.5 Let $T : D \subset X \rightarrow X$ be an operator on a metric space (X, d) and let $\mu \in D$. Suppose that

$$d(T(x), \mu) \leq \varphi(d(x, \mu)) \text{ for each } x \in D \text{ with } d(x, \mu) \in I,$$

where φ is a strict gauge function of order $\delta \geq 1$ on I . Then, μ is a unique fixed point of T in the set $U = \{x \in D : d(x, \mu) \in I\}$. Moreover, if $T : U \rightarrow U$, then for each $x_0 \in U$ the following items hold.

- (i) Picard iteration $x_{n+1} = T(x_n)$ stays in U and converges to μ with Q -order δ .
- (ii) $d(x_n, \mu) \leq \lambda^{s_n(\delta)} d(x_0, \mu)$ for each $n = 0, 1, 2, \dots$ where $\lambda = \phi(E(x_0))$ and ϕ is a non-decreasing nonnegative function on I satisfying $\varphi(t) = t\phi(t)$ for each $t \in I$.
- (iii) $d(x_{n+1}, \mu) \leq \varphi(d(x_n, \mu))$ for each $n = 0, 1, 2, \dots$

3 Local convergence analysis

The local convergence analysis of Newton’s method is based on generalized w - affine invariant conditions. Let p be such that $F(p) = 0$ and $F'(p)^{-1}$ exists. Suppose there exists function $w_0 : [0, +\infty) \rightarrow [0, +\infty)$ continuous and increasing with $w_0(0) = 0$ such that for each $x \in D$

$$\|F'(p)^{-1}(F'(x) - F'(p))\| \leq w_0(\|x - p\|). \quad (3.1)$$

Moreover, suppose equation

$$w_0(t) = 1 \quad (3.2)$$

has positive solutions. Denote by ρ the smallest positive solution of equation (3.2). Set $D_0 = D \cap U(x_0, \rho)$. Furthermore, suppose there exists function $w : [0, 2\rho) \times [0, \rho) \times [0, \rho) \rightarrow [0, +\infty)$ continuous and nondecreasing with $w(0, 0, 0) = 0$ such that for each $x, y \in D_0$

$$\|F'(p)^{-1}(F'(x) - F'(y))\| \leq w(\|x - y\|, \|x - p\|, \|y - p\|). \quad (3.3)$$

Notice that function w_0 depends on D , whereas w depends on w_0 and D_0 . The construction of function w was not possible before, since w_0 and D_0 are needed [3, 9, 10, 13, 22, 33]:

$$\begin{aligned} & \|F'(p)^{-1}(F'(x) - F'(y))\| \\ & \leq \bar{w}(\|x - y\|, \|x - p\|, \|y - p\|) \quad (3.4) \\ & , \text{ for each } x, y \in D \end{aligned}$$

where $\bar{w} : [0, +\infty)^3 \rightarrow [0, +\infty)$ is a continuous and nondecreasing function with $\bar{w}(0, 0, 0) = 0$. We have that for $s, s_1, s_2, s_3 \geq 0$

$$w_0(s) \leq \bar{w}(s, s, s) \quad (3.5)$$

and

$$w(s_1, s_2, s_3) \leq \bar{w}(s_1, s_2, s_3) \quad (3.6)$$

since $D_0 \subseteq D$ and $\frac{\bar{w}}{w_0}$ can be arbitrarily large [3, 4]. Let us provide a motivational example to show that (3.5) and (3.6) can hold as strict inequalities. Let $\mathcal{B}_1 =$

$\mathcal{B}_2 = \mathbb{R}^3, D = \bar{U}(0, 1)$ and $p = (0, 0, 0)^T$. Define function F on D for $v = (v_1, v_2, v_3)^T$ by

$$F(v) = (e^{v_1} - 1, \frac{e-1}{2}v_2^2 + v_2, v_3)^T. \quad (3.7)$$

The Fréchet derivative is given by

$$F'(v) = \begin{bmatrix} e^{v_1} & 0 & 0 \\ 0 & (e-1)v_2 + 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Then, conditions (3.1), (3.3) and (3.4) are satisfied for $w_0(t) = L_0t, w(t, t, t) = e^{\frac{1}{L_0}t}, \bar{w}(t, t, t) = Lt, \rho = \frac{1}{L_0}, L_0 = e - 1$ and $L = e$. Notice that

$$w_0(t) < w(t, t, t) < \bar{w}(t, t, t) \quad (3.8)$$

holds for all $t \in [0, \rho]$. The upper bound on the expression appearing in these studies using (3.4) is

$$\begin{aligned} & \|F'(p)^{-1}(F(x) - F(p) - F'(x)(x - p))\| \\ & \leq \left\| \int_0^1 F'(p)^{-1}(F'(x) - F'(p + \theta(x - p))) \right. \\ & \quad \times (x - p) d\theta \left. \right\| \\ & \leq \int_0^1 \bar{w}(\|x - p\|, \theta\|x - p\|, (1 - \theta)\|x - p\|) d\theta \\ & = \int_0^{\|x-p\|} \bar{w}(\|x - p\|, u, \|x - p\| - u) du \quad (3.9) \end{aligned}$$

However, using the more precise and actually needed condition (3.3) instead of (3.4) we obtain instead of (3.9) the more precise estimate (see also (3.6))

$$\begin{aligned} & \|F'(p)^{-1}(F(x) - F(p) - F'(x)(x - p))\| \\ & \leq \left\| \int_0^1 F'(p)^{-1}(F'(x) - F'(p + \theta(x - p))) \right. \\ & \quad \times (x - p) d\theta \left. \right\| \\ & \leq \int_0^1 w(\|x - p\|, \theta\|x - p\|, (1 - \theta)\|x - p\|) d\theta \\ & = \int_0^{\|x-p\|} w(\|x - p\|, u, \|x - p\| - u) du. \quad (3.10) \end{aligned}$$

Replacing (3.4) and (3.9) by (3.3) and (3.10), respectively in the proof of the local convergence analysis of Newton’s method in [33] we obtain a finer convergence analysis with advantages as already stated in the introduction of this study. The advantages are obtained under the same computational effort as in [33], since in practice the computation of function \bar{w} requires the computation of function w as a special case.

The results in this Section, in particular reduce to the corresponding ones in Section 7 of [33], if $w_0 = w = \bar{w}$. Otherwise, the new results constitute an improvement (see also advantages (A)).

LEMMA 3.1 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open convex set D . Let $p \in D$ be a simple zero of F and the condition (3.1) and (3.3) are satisfied. Assume that the real function

$$\varphi(t) = \frac{\int_0^1 w(t, u, t - u) du}{1 - w_0(t)} \quad (3.11)$$

is well-defined on the interval $(0, \rho)$. Then

$$\|N(x) - p\| \leq \varphi(\|x - p\|) \text{ for each } x \in D \text{ with } \|x - p\| < \rho. \quad (3.12)$$

Proof. Let $x \in D$ be such that $\|x - p\| < \rho$. Notice that since function φ is well-defined on $[0, \rho)$, then $w_0(t) < 1$ for each $t \in [0, \rho)$. Using this and (3.1), we get in turn that

$$\|F'(p)^{-1}(F'(x) - F'(p))\| \leq w_0(\|x - p\|) < 1,$$

since $0 \leq \|x - p\| < \rho$. It then follows from the preceding inequality and the Banach Lemma on invertible operators [5, 22] that $F'(x)^{-1}$ exists and

$$\|F'(x)^{-1}F'(p)\| \leq \frac{1}{1 - w_0(\|x - p\|)}.$$

Moreover, we have that

$$\begin{aligned} & \|T(x) - p\| \\ & \leq \|F'(x)^{-1}F'(p)\| \\ & \quad \times \left\| \int_0^1 F'(p)^{-1}(F'(x) - F'(p + \theta(x - p))) \right. \\ & \quad \times (x - p) d\theta \left. \right\| \\ & \leq \|F'(x)^{-1}F'(p)\| \\ & \quad \times \int_0^1 w(\|x - p\|, \theta\|x - p\|, \\ & \quad \|x - (p + \theta(x - p))\|) d\theta \\ & \leq \|F'(x)^{-1}F'(p)\| \\ & \quad \times \int_0^{\|x-p\|} w(\|x - p\|, u, \|x - p\| - u) d\theta. \end{aligned}$$

The proof is completed if we combine the last two inequalities. □

THEOREM 3.2 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose F' satisfies conditions (3.1) and (3.3) and the real function φ defined by (3.11) is strict gauge function of order $\xi + 1$ for some $\xi \geq 0$ on the interval $[0, \rho)$. Then for each $x_0 \in U(p, \rho)$ the following items hold true:

- (i) Newton iteration (1.1) is well-defined, remains in $U(p, \rho)$ and converges to p with Q -order $\xi + 1$.
- (ii) For all $n \geq 0$ we have the following estimate

$$\|x_{n+1} - p\| \leq \varphi(\|x_n - p\|). \quad (3.13)$$

- (iii) For all $n \geq 0$ we have the following estimate

$$\|x_n - p\| \leq \lambda^{s_n(p+1)} \|x_0 - p\|, \quad (3.14)$$

where $\lambda = \phi(\|x_0 - p\|)$ and ϕ is a nondecreasing function on I satisfying $\varphi(t) = t\phi(t)$.

- (iv) If ρ is a fixed point of φ , then ρ is the optimal radius of the convergence ball of Newton's method under the condition (3.1) and (3.3) for some w and w_0 .

Proof. Items (i)-(iii) follow from Corollary 2.5 and Lemma 3.1. Item (iv) is shown in Theorem 4.1. □

4 Special cases

We assume from now on that

$$w_0(t) \leq w_1(t) \text{ for } t \in [0, \rho]. \quad (4.1)$$

If

$$w_1(t) \leq w_0(t) \text{ for } t \in [0, \rho], \quad (4.2)$$

where w_1 is a nonnegative nondecreasing function of $[0, \rho]$. Then, the results that follow hold with w_0 replacing w_1 . In the following theorems and corollaries in this section we consider some interesting special cases of Theorem 3.2.

THEOREM 4.1 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose

$$\|F'(p)^{-1}(F'(x) - F'(y))\| \leq w_1(\|x - p\|) - w_1(\|y - p\|) \quad (4.3)$$

for all $x, y \in U(p, \rho)$, where w_1 is a real function defined on $[0, \rho]$ with $w_1(0) = 0$. Moreover, suppose that (3.1) holds. Furthermore, suppose that

$$\varphi(t) = \frac{tw_1(t) - \int_0^1 w_1(u)du}{1 - w_0(t)} \quad (4.4)$$

is a strict gauge function of order $\xi + 1$ for some $p \geq 0$ on $[0, r)$. Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -order $\xi + 1$ to p and satisfies the estimates (4.1) and (4.4). Moreover, if ρ is a fixed point of φ and w_0, w_1 are continuous, then ρ is the optimal radius of the convergence ball of Newton's method.

Proof. The first part of the Theorem follows immediately from Theorem 3.2. Let ρ be a fixed point of φ . We shall show the exactness of ρ even if $\mathcal{B}_1 = \mathcal{B}_2 = \mathbb{R}$. Let $p \in \mathbb{R}_+$ be arbitrary. Define function F on $D = \bar{U}(p, \rho)$ by

$$F(x) = x - p - \text{sign}(x - p) \int_0^{\|x - p\|} w_1(t)dt,$$

where w_1 is continuous on $[0, \rho]$ with $w_1(0) = 0$. Clearly, F is continuously differentiable with $F'(x) = 1 - w_1(|x - p|)$. Then, for all $x \in U(p, \rho)$ and $y \in [p, x]$, we get

$$\|F'(p)^{-1}(F'(x) - F'(y))\| = w_1(|x - p|) - w_1(|y - p|)$$

which shows that (4.3) holds. Then, $T(p + \rho) = p - \varphi(\rho) = p - \rho$ and $T(p - \rho) = p + \varphi(\rho) = p + \rho$, for $T(x) = x - F'(x)^{-1}F(x)$. Hence, if $x_0 = p + \rho$, then $x_n = p + (-)^n \rho$. Therefore, Newton's method starting at $x_0 = p + \rho$ fails to converge. □

REMARK 4.2 Let us give a sufficient condition for φ defined by (4.4) to be a gauge function of order $\xi + 1$. It follows from the Example in section 3 that if w_1 is a nonnegative nondecreasing function on $[0, \rho]$ such that for all $\lambda \in (0, 1)$ and all $t, u \in [0, \rho]$ with $t \geq u$ it satisfies $w_1(\lambda t) - w_1(\lambda u) \leq \lambda^\xi [w_1(t) - w_1(u)]$ for some $\xi \geq 0$, then the function φ defined by (4.4) is a strict gauge function of order $\xi + 1$ on I provided that $\varphi(t) < t$ for all $t \in (0, \rho)$.

Note that in the case $w_0(t) = w_1(t) = Lt$ condition (4.3) coincides with (4.2) and we obtain Traub and Woźniakowski's result [39]. By putting $w_1(t) = \int_0^t L(u)du$, where L is nondecreasing on $[0, \rho)$, we immediately get some results of Wang [43, Theorem 3.1 and 5.1]. Theorem 4.1 is also an improvement of a result by Wang and Li [45, Theorem 1.1] and Proinov [33].

COROLLARY 4.3 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose (4.3) holds with

$$w_0(t) = L_0 t^\xi, \quad (4.5)$$

$$w_1(t) = L_1 t^\xi \quad (\xi > 0, L_0, L_1 > 0) \text{ and}$$

$$0 < r \leq R = \left(\frac{\xi + 1}{\xi L_1 + (\xi + 1)L_0} \right)^{1/\xi}.$$

Define the real functions

$$\varphi(t) = \frac{\xi}{\xi + 1} \frac{L_1 t^{\xi+1}}{1 - L_0 t^\xi} \text{ and } \phi(t) = \frac{\xi}{\xi + 1} \frac{L_1 t^\xi}{1 - L_0 t^\xi}. \quad (4.6)$$

Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -order $\xi + 1$ to p and satisfies the estimates (4.1) and (4.4). Moreover, if $\rho = R$ is the optimal radius of the convergence ball of Newton's method under the condition (4.3) with w_1 defined by (4.5).

In the case when $\xi = 1$ and $L_0 = L_1$, we again get the above mentioned result of Traub and Woźniakowski [39] as well as the result of Rheinboldt [35], Wang [42] and Ypma [49]. If $0 < \xi \leq 1$, then Corollary 4.3 leads to the results of Wang and Li [45, Corollary 3.1], Huang [19, Theorem 2] and Proinov [33]. If $L_0 < L_1$ the mentioned results are improved.

COROLLARY 4.4 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose that (3.1) and (4.3) hold with

$$\begin{aligned} w_0(t) &= \frac{c_0}{(1 - \gamma_0 t)^2}, & (4.7) \\ w_1(t) &= \frac{c}{(1 - \gamma t)^2}, (\gamma > 0, L_0, L_1 > 0) \text{ with} \\ c_0 &\leq c \text{ and } \gamma_0 \leq \gamma \end{aligned}$$

Define the real function

$$\Phi(t) = \frac{\varphi(t)}{t} \tag{4.8}$$

and

$$0 < r \leq R, \tag{4.9}$$

where R is the smallest positive solution of $\varphi(t) = t$. Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -quadratically to p with the estimates (4.1) and

$$\|x_n - p\| \leq \lambda^{2^n - 1} \|x_0 - p\| \text{ for all } n \geq 0, \tag{4.10}$$

where $\lambda = \Phi(\|x_0 - p\|)$. Moreover, if $\rho = R$ is the optimal radius of the convergence ball of Newton's method under the conditions (3.1) and (4.3).

Corollary 4.4 without the estimate (4.1) and for $L_0 = L, \gamma_0 = \gamma$ is due to Wang and Han [44] ($L = 1$) and Wang [43, Example 1] ($L \geq 0$). Note that the function Φ defined in (4.8) is strictly increasing and continuous on the interval $I = [0, R]$ and satisfies $\Phi(I) = [0, 1]$. Therefore, if we ignore the estimate (4.1), then Corollary 4.4 is equivalent to the following result of Wang [43, p.132]. However, if $L_0 < L$ and $\gamma_0 < \gamma$ the mentioned results are improved.

COROLLARY 4.5 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose that (3.1) and (4.3) hold and R satisfies (4.9). Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -quadratically to p with the estimates (4.1) and

$$\|x_n - p\| \leq \lambda^{2^n - 1} \|x_0 - p\| \text{ for all } n \geq 0. \tag{4.11}$$

The following theorem is another natural generalization of Traub and Woźniakowski's result [39] mentioned in the beginning of the section.

THEOREM 4.6 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose that (3.1) and

$$\|F'(p)^{-1}(F'(x) - F'(y))\| \leq w_2(\|x - y\|) \tag{4.12}$$

for all $x, y \in U(p, \rho)$ where w_2 is a real function defined on $[0, \rho]$ with $w_2(0) = 0$. Assume that

$$\varphi(t) = \frac{\int_0^t w_2(u) du}{1 - w_0(t)} \tag{4.13}$$

is a strict gauge function of order $\xi + 1$ for some $0 \leq \xi \leq 1$ on $[0, \rho)$. Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -order $\xi + 1$ to p and with the error estimates (4.1) and (4.5).

COROLLARY 4.7 Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose conditions (3.1) and (4.12) hold with

$$\begin{aligned} w_0(t) &= L_0 t^\xi, & (4.14) \\ w_2(t) &= L_2 t^\xi, (0 \leq \xi \leq 1, L_0, L_2 > 0) \text{ and} \\ 0 < r &\leq R = \left(\frac{\xi + 1}{\xi L_2 + (\xi + 1)L_0} \right)^{1/\xi}. \end{aligned}$$

In the case $\xi = 0$ we assume that $L_0 < \frac{1}{2}$ and $R = \infty$. Define functions

$$\varphi(t) = \frac{\xi}{\xi + 1} \frac{L_2 t^{\xi+1}}{1 - L_0 t^\xi} \text{ and } \phi(t) = \frac{\xi}{\xi + 1} \frac{L_2 t^\xi}{1 - L_0 t^\xi}. \tag{4.15}$$

Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -order $\xi + 1$ to p and satisfies the estimates (4.1) and (4.4).

From Corollary 4.7 in the case $\xi = 1$ we again get the classical results of Rheinboldt [35, 36], Traub and Woźniakowski [39], Wang [42–45], Ypma [49]. In the case $0 \leq \xi \leq 1$ Corollary 4.7 is obtained by Ypma [49, Theorem 3.1], Huang [19, Theorem 1] and in a slightly different form by Argyros [6, Theorem 4].

THEOREM 4.8 *Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose that (3.1) and*

$$\|F'(p)^{-1}(F'(x) - F'(y))\| \leq w_3(\|x - y\|) \quad (4.16)$$

for all $x, y \in U(p, \rho)$, where w_3 is a real function defined on $[0, \rho]$ with $w_3(0) = 0$. Assume that

$$\varphi(t) = \frac{tw_3(t) + \int_0^t w_3(u)du}{1 - w_0(t)} \quad (4.17)$$

is a strict gauge function of order $\xi + 1$ for some $0 \leq \xi \leq 1$ on $[0, \rho)$. Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -order $\xi + 1$ to p and with the error estimates (4.1) and (4.5).

REMARK 4.9 *Let φ be defined by (4.15) or (4.18). It follows from the Example 2.2 in [33] that if w_3 is nonnegative nondecreasing on $[0, \rho)$ and $\frac{w_3(t)}{t^\xi}$ is nondecreasing on $(0, \rho)$ for some $\xi \geq 0$, then φ is a strict gauge function of order $\xi + 1$ on $[0, \rho)$ provided that $\varphi(t) < t$ for all $t \in (0, \rho)$.*

COROLLARY 4.10 *Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose that (4.16) hold with*

$$\begin{aligned} w_0(t) &= L_0 t^\xi, \\ w_3(t) &= L_3 t^\xi, (\xi > 0, L_0 > 0) \text{ and} \\ 0 < \rho \leq R &= \left(\frac{\xi + 1}{\xi L_3 + (\xi + 1)L_0} \right)^{1/\xi} \end{aligned} \quad (4.18)$$

In case $\xi = 0$ we assume That $L_0 < \frac{1}{3}$ and $R = \infty$. Define the real functions

$$\varphi(t) = \frac{(\xi + 2)L_3 t^{\xi+1}}{(\xi + 1)(1 - L_0 t^\xi)} \text{ and } \phi(t) = \frac{(\xi + 2)L_3 t^\xi}{(\xi + 1)(1 - L_0 t^\xi)}. \quad (4.19)$$

Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -quadratically to p with the estimates (4.3) and (4.5).

From Corollary 4.10 in the case $\xi = 1$ we obtain $R = \frac{2}{5L}$ which improves a recent result of Wang and Li [45, Corollary 3.2]. For example they have proved $R \leq \frac{1}{3L_0}$, since Proinov [33] $R_\xi = \frac{2}{5L} < \frac{2}{5L_0}$, The following corollary is an improvement of another result of Wang and Li [45, Corollary 3.3].

COROLLARY 4.11 *Let $F : D \subset \mathcal{B}_1 \rightarrow \mathcal{B}_2$ be a continuously Fréchet differentiable operator on an open ball $U(p, \rho) \subset D$, where p is a simple zero of F . Suppose that (4.16) holds with w_3 defined by (4.18) and R satisfies (4.18). Then starting from every $x_0 \in U(p, \rho)$ Newton iteration (1.1) is well defined, remains in $U(p, \rho)$, converges with Q -quadratically to p with the estimates (4.3) and (4.5).*

In the convergence ball of Newton’s method, the solution p of the equation $F(x) = 0$ is certainly unique. But it is well known that the uniqueness ball of this equation may be larger. Such results can be found in [6, 33] using only (3.1).

EXAMPLE 4.12 *Returning back at the motivational example (3.7), we see by Lemma 3.1 and Theorem 3.2 that our new radius of convergence is $\rho_3 = \frac{2}{2(e-1)+e^{\frac{1}{e-1}}} = 0.3826919122323857$ the radius in [3, 4, 6] is $\rho_2 = \frac{2}{2(e-1)+e} = 0.324947231372689$ and the one in [19, 33, 39, 41] is $\rho_1 = \frac{2}{3e} = 0.2452529607809$, so $\rho_1 < \rho_2 < \rho_3$ justifying advantages (A).*

EXAMPLE 4.13 *Let $\mathcal{B}_1 = \mathcal{B}_2 = C[0, 1]$, the space of continuous functions defined on $[0, 1]$ equipped with the max norm and $D = \bar{U}(0, 1)$. Define the operator F on D by*

$$F(x)(t) = x(t) - 5 \int_0^1 t\theta x^3(\theta)d\theta,$$

so

$$F'(x)u(t) = u(t) - 15 \int_0^1 t\theta x^2(\theta)u(\theta)d\theta \text{ for each } u \in D.$$

Then, we have for $p(t) = 0$ ($t \in [0, 1]$), $\xi = 1$, that $L_0 = 7.5, L_3 = L = 15, w_0(t) = 7.5t, w_3(t) = 15t$. Using (4.18), the radius in [19, 33, 39, 41] is $\rho_1 = \frac{2}{3L_3} = 0.0444$ whereas the new radius is $\rho_3 = \frac{2}{2L_0+L_3} = 0.0667$. Hence, our radius of convergence is larger.

5 Conclusion

The aim of this paper is to provide a finer local convergence analysis for Newton’s method than in earlier

papers using the gauge theory, the w -conditions, the center w_0 -conditions and our new idea of restricted convergence regions. Using the center w_0 -condition instead of the less precise w -condition used earlier, we find tighter upper bounds on the $\|F(x)^{-1}F(x_0)\|$. Moreover, using restricted convergence regions, we obtain a more precise location where the Newton iterates lie leading to tighter w -majorants. This way, we obtain a larger radius of convergence resulting to a wider choice of initial guesses and tighter error estimates on the distances $\|x_n - p\|$ leading to the calculation of fewer iterates to obtain a desired error tolerance. Moreover, the information on the location of the solution is also more accurate than in earlier studies. These, improvements are made under the same computational cost, since in practice the computation of old w -functions requires the computation of the new w_0 - and w_1 functions as special cases.

References:

- [1] S.Amat S. Busquier and S.Plaza , *Dynamics of the King and Jarratt iterations* , Aequationes Math. 69(3), (2005), 212–223 .
- [2] S.Amat S. Busquier and S.Plaza , *Chaotic dynamics of a third-order Newton-type method*, J. Math. Anal. Appl. 366(1),(2010) pp. 24–32.
- [3] I.K. Argyros, On the convergence and application of Newton's method under weak Hölder continuity assumptions, Int. J. Comput. Math. 80 (2003), 767-780.
- [4] I.K. Argyros, J.M. Gutiérrez, A unified approach for enlarging the radius of convergence for Newton's method and applications, Nonlinear Funct. Anal. Appl. 10 (2005), 555-563.
- [5] I. K. Argyros and A. Magreñán, Iterative methods and their dynamics with applications, CRC Press, New York, 2007.
- [6] I. K. Argyros and S. Hillout, Weaker conditions for the convergence of Newton's method. J. Complexity 28, 364–387, 2012.
- [7] L. Blum, F. Cucker, M. Shub, S. Smale, Complexity and Real Computation, Springer, New York, 1998.
- [8] F. Cianciaruso, E. De Pascale, Newton-Kantorovich approximations when the derivative is Hölderian: old and new results, Numer. Funct. Anal. Optim. 24 (2003), 713-723.
- [9] F. Cianciaruso, E. De Pascale, Estimates of majorizing sequences in the Newton-Kantorovich method, Numer. Funct. Anal. Optim. 27 (2006), 529-538.
- [10] F. Cianciaruso, E. De Pascale, Estimates of majorizing sequences in the Newton-Kantorovich method: A further improvement, J. Math. Anal. Appl. 332 (2006), 329-335.
- [11] E. De Pascale, P.P. Zabrejko, The convergence of the Newton-Kantorovich method under Verweim conditions: a new improvement, Z. Anal. Anwend. 17 (1998), 271-280.
- [12] J.-P. Dedieu, Condition number analysis for sparse polynomial systems, in: F. Cucker, M. Shub (Eds.), Foundations of Computational Mathematics, Springer, Berlin, 1997, 75-101.
- [13] K. Dočev, Über Newtonsche Iterationen, C. R. Acad. Bulg. Sci. 36 (1962), 695-701.
- [14] J. A. Ezquerro, M. A. Hernández, Generalized differentiability conditions for Newton's method, IMA J. Numer. Anal. 22 (2002), 187-205.
- [15] J. A. Ezquerro, M. A. Hernández, On the R -order of convergence of Newton's method under mild differentiability conditions, J. Comput. Appl. Math. 197 (2006), 53-61.
- [16] A. Galperin, On Convergence domain of Newton's and modified Newton's methods, Numer. Funct. Anal. Optim. 26 (2005), 385-405.
- [17] W. B. Gragg, R. A. Tapia, Optimal error bounds for the Newton-Kantorovich theorem, SIAM J. Numer. Anal. 11 (1974), 10-13.
- [18] M. A. Hernández, The Newton method for operators with Hölder continuous first derivative, J. Optimization Theory Appl. 109 (2001), 631-648.
- [19] Z.D. Huang, The convergence ball of Newton's method and uniqueness ball of equations under Hölder-type continuous derivatives, Comput. Math. Appl. 47 (2004), 247-251.
- [20] L.O. Jay, A note on Q-order of convergence, BIT 41 (2001), 422-429.
- [21] B. Jankó, Rezolvarea equatiilor operationale neliniare in spatii Banach, Editura Academiei republicii socialiste Romania, Bucuresti, 1969.

- [22] L.V. Kantorovich, G.P. Akilov, *Functional Analysis in Normed Spaces*, Pergamon, Oxford, 1982.
- [23] H. B. Keller, Newton's methods under mild differentiability conditions, *J. Comput. System Sci.* 4 (1970), 15-28.
- [24] Yu. V. Lysenko, The convergence conditions of the Newton-Kantorovich method for nonlinear equations with Hölder linearizations (Russian), *Dokl. Akad. Nauk BSSR* 38 (1994), 20-24.
- [25] A. A. Magreñán, Different anomalies in a Jarratt family of iterative root finding methods, *Appl. Math. Comput.* 233, (2014), 29-38.
- [26] A. A. Magreñán, A new tool to study real dynamics: The convergence plane, *Appl. Math. Comput.* 248, (2014), 29-38.
- [27] G. J. Miel, An update version of the Kantorovich theorem for Newton's method, *Computing* 27 (1981), 237-244.
- [28] A. M. Ostrowski, La méthode de Newton dans les espaces de Banach, *C. R. Acad. Sci. Paris sér A* 272 (1971), 1251-1253.
- [29] D. Petcu, On the Kantorovich hypothesis for Newton's method, *Informatika (Vilnius)* 4 (1993) No. 1-2, 188-198.
- [30] F. A. Potra, On the a posteriori error estimates for Newton's method, *Beitraege Numer. Math.* 12 (1984), 125-138.
- [31] P.D. Proinov, A generalization of the Banach contraction principle with high order of convergence of successive approximations, *Nonlinear Anal.* 67 (2007), 2361-2369.
- [32] P.D. Proinov, New general convergence theory for iterative processes and its applications to Newton-Kantorovich-type theorems, *J. Complexity*, 25, (2009), 38-62.
- [33] P.D. Proinov, General local convergence theory for a class of iterative processes and its applications to Newton's method, *J. Complexity*, 25, (2009), 38-62.
- [34] L.B. Rall, A note on the convergence of Newton's method, *SIAM J. Numer. Anal.* 11 (1974), 34-36.
- [35] W.C. Rheinboldt, An adaptive continuation process for solving systems of nonlinear equations, in: *Banach center publications*, Vol. 3, Polish Academy of science, Warsaw, 1977, 129-142.
- [36] W. C. Rheinboldt, On a theorem of S. Smale about Newton's method for analytic mappings, *Appl. Math. Lett.* 1 (1988), 69-72.
- [37] E. Schröder, Über unendlich viele Algorithmen zur Auflösung der Gleichungen, *Math. Anal.* 2 (1870), 317-365.
- [38] S. Smale, Newton's method estimates from data at one point, in: R.E. Ewing, K.E. Gross, C.F. Martin (Eds.), *The Merging of Disciplines: New Direction in Pure, Applied, and Computational Mathematics*, Springer, New York, 1986, 185-196.
- [39] J.F. Traub, H. Wozniakowski, Convergence and complexity of Newton iteration for operator equations, *J. Assoc. Comput. Mach.* 26 (1979), 250-258.
- [40] B. A. Vertgeim, On the Conditions for the applicability of Newton's method (Russian), *Dokl. Akad. Nauk SSSR* 110 (1956), 719-722.
- [41] B. A. Vertgeim, On some methods of the approximate solution of nonlinear functional equations in Banach spaces (Russian), *Uspekhi Mat. Nauk* 12 (1957), 166-169 [Engl. Transl: *Amer. Math. Soc. Transl.* 16 (1960), 378-382].
- [42] X.H. Wang, The convergence of Newton's method, *Kexue Tongbao* 25 (1980), 36-37 (in Chinese).
- [43] X.H. Wang, Convergence of Newton's method and uniqueness of the solution of the equations in Banach space, *IMA J. Numer. Anal.* 20 (2000), 123-134.
- [44] X.H. Wang, D.F. Han, Criterion α and Newton's method under weak conditions, *Math. Numer. Sin.* 19 (1997) 103-112 (in Chinese) English translation in: *Chinese J. Numer. Math. Appl.* 19 (1997), 96-105.
- [45] X.H. Wang, C. Li, Convergence of Newton's method and uniqueness of the solution of the equations in Banach space. II, *Acta. Math. Sinica (Engl. Ser.)* 19 (2003), 405-412.
- [46] T. Yamamoto, A unified derivation of several error bounds for Newton's process, *J. Comput. Appl. Math.* 12-13 (1985), 179-191.
- [47] T. Yamamoto, Historical development in convergence analysis for Newton's and Newton-like methods, *J. Comput. Appl. Math.* 124 (2000), 1-23.

- [48] J.-C. Yakoubsohn, Finding a cluster of zeros of univariate polynomials, *J. Complexity* 16 (2000), 603-638.
- [49] T.J. Ypma, Local convergence of inexact Newton methods, *SIAM J. Numer. Anal.* 21 (1984), 583-589.