

# On a Class of Multivalent Functions With Negative Coefficients Involving $(r, q)$ – Calculus

MA'MOUN I. Y. ALHARAYZEH<sup>1\*</sup>, MASLINA DARUS<sup>2</sup>, HABIS S. AL-ZBOON<sup>3</sup>

<sup>1</sup>Department of Scientific Basic Sciences, Faculty of Engineering Technology  
Al-Balqa Applied University, Amman 11134  
JORDAN

<sup>2</sup>Department of Mathematical Sciences, Faculty of Science and Technology  
Universiti Kebangsaan Malaysia, Bangi, 43600 Selangor D. Ehsan,  
MALAYSIA

<sup>3</sup>Department of Curriculum and Instruction, College of Education  
Al-Hussein Bin Talal University,  
JORDAN

*Abstract:* - In this research, we focused on presenting a novel subclass of multivalent analytic functions situated in the open unit disk, characterized by the use of Jackson's derivative operator. Our investigation systematically establishes the requisite inclusion conditions in this class, offering detailed coefficient characterizations. The exploration encompasses an array of significant properties intrinsic to this subclass, encompassing coefficient estimates, growth and distortion theorems, identification of extreme points, and the determination of the radius of starlikeness and convexity for functions falling within this specialized category. Expanding the preliminary findings, this research extended the inquiry to delve deeper into the intriguing features and implications associated with this new subclass of multivalent analytic functions. The research concentrated the light on the nuanced intricacies of coefficient estimates, providing a comprehensive understanding of how these functions evolve within the open unit disk through exploring the growth and distortion theorems, unraveling the underlying mathematical principles governing the behavior of functions in this subclass as they extend beyond the unit disk. The findings of this research contribute to the broader understanding of multivalent analytic functions, paving the way for further exploration and applications in diverse mathematical contexts.

*Key-Words:* -Analytic function, Unit disk, New subclass,  $p$ -valent function, Quantum or  $(r, q)$ -Calculus,  $(r, q)$ -Derivative operator.

Received: October 24, 2023. Revised: February 25, 2024. Accepted: March 13, 2024. Published: April 10, 2024.

## 1 Commencement and Definition

The category of all analytic functions exhibiting the following structure

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad a_n \text{ is complex number}$$

within the open unit disk, denoted as  $\mathcal{U} = \{z \in \mathbb{C} : |z| < 1\}$  where  $\mathbb{C}$  signifies the set of complex numbers. This class of all analytic functions is denoted as  $\hat{\mathcal{A}}$ . Also, let  $\hat{\mathcal{A}}(p) (p \in \mathbb{N} = \{1, 2, 3, \dots\})$  be the class consisting of all analytic functions  $f$ . This class is represented by a series that articulates the underlying structure of these functions.

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n, \quad a_n \text{ is complex number}$$

which called  $p$ -valent in the open unit disk  $\mathcal{U}$  over the complex numbers  $\mathbb{C}$ . It is important to observe that  $\hat{\mathcal{A}}(1)$  is equivalent to  $\hat{\mathcal{A}}$ . Additionally, the collection of all univalent functions within the open disk  $\mathcal{U}$  is symbolized as  $S(p)$  which is a subclass of  $\hat{\mathcal{A}}(p)$ . Furthermore, let  $S_p^*(\alpha)$  and  $C_p(\alpha)$  represent the classes of  $p$ -valent functions respectively starlike of order  $\alpha$  and convex of order  $\alpha$ , for  $0 \leq \alpha < p$ . Notably,  $S_p^*(0)$  is synonymous with  $S_p^*$  and  $C_p(0)$  corresponds to  $C_p$ , both being the well-known classes of starlike and convex  $p$ -valent functions in  $\mathcal{U}$ , respectively.

Next, let us assume that  $\mathcal{T}(p) (p \in \mathbb{N} = \{1, 2, 3, \dots\})$  denote the subclass of  $S(p)$  of analytic functions

having the structure

$$f(z) = z^p - \sum_{n=p+1}^{\infty} a_n z^n, \quad a_n > 0. \quad (1)$$

As it is defined on the open unit disk  $\mathcal{U} = \{z \in \mathbb{C} : |z| < 1\}$ . A function  $f \in \mathcal{T}(p)$  is denoted as a  $p$ -valent function with negative coefficients. Moreover, it is evident that  $S_{\mathcal{T},p}^*(\alpha)$  and  $C_{\mathcal{T},p}(\alpha)$  for  $0 \leq \alpha < p$ , that are  $p$ -valent functions, respectively starlike of order  $\alpha$  and convex of order  $\alpha$  which is subclasses of  $\mathcal{T}(p)$ . Clearly, the class  $\mathcal{T}(1) = \mathcal{T}$  in [1], he derived and investigated the subclasses of  $\mathcal{T}(1)$  denoted by  $S_{\mathcal{T},1}^*(\alpha) = S_{\mathcal{T}}^*(\alpha)$  and  $C_{\mathcal{T},1}(\alpha) = C_{\mathcal{T}}(\alpha)$ , for  $0 \leq \alpha < 1$  that are respectively starlike of order  $\alpha$  and convex of order  $\alpha$ .

In this section, we revisit established concepts and fundamental results of  $(r, q)$ -calculus. Throughout this paper, we denote constants as let  $r, q$  be constants with  $0 < q < r \leq 1$ . We provide some definitions and theorems pertinent to  $(r, q)$ -calculus, which will be employed in the subsequent sections of these papers. [2], [3], [4], [5], [6], and, [7].

For  $0 < q < r \leq 1$  the Jackson's  $(r, q)$ -derivative of a function  $f \in \hat{\mathcal{A}}(p)$  is, by definition, given as follow

$$\mathcal{D}_{r,q}f(z) := \begin{cases} \frac{f(rz) - f(qz)}{(r-q)z}, & z \neq 0, \\ f'(0), & z = 0. \end{cases} \quad (2)$$

From (2), we have

$$\mathcal{D}_{r,q}f(z) = [p]_{r,q} z^{p-1} + \sum_{n=p+1}^{\infty} [n]_{r,q} a_n z^{n-1},$$

where  $[p]_{r,q} = \frac{r^p - q^p}{r - q}$ ,  $[n]_{r,q} = \frac{r^n - q^n}{r - q}$  and  $0 < q < r \leq 1$ .

Note that for  $r = 1$ , the Jackson  $(r, q)$ -derivative reduces to the Jackson  $q$ -derivative operator of the function  $f$ ,  $\mathcal{D}_q f(z)$  (refer to [8], [9], and, [10]). Note also that  $\mathcal{D}_{1,q} f(z) \rightarrow f'(z)$  when  $q \rightarrow 1^-$ , where  $f'$  is the classical derivative of the function  $f$ .

Clearly for a function  $g(z) = z^n$ , we obtain

$$\mathcal{D}_{r,q}g(z) = \mathcal{D}_{r,q}z^n = \frac{r^n - q^n}{r - q} z^{n-1} = [n]_{r,q} z^{n-1}.$$

And

$$\lim_{q \rightarrow 1^-} \mathcal{D}_{1,q}g(z) = \lim_{q \rightarrow 1^-} \frac{1 - q^n}{1 - q} z^{n-1} = n z^{n-1} = g'(z),$$

where  $g'$  is the ordinary derivative.

The theory of  $q$ -calculus finds application in the adaptation and resolution of various problems in applied science like ordinary fractional calculus, quantum physics, optimal control, hypergeometric series, operator theory,  $q$ -difference and  $q$ -integral equations, as well as geometric function theory of complex analysis. The application of  $q$ -calculus was started by, [11]. In, [12], have utilized the fractional  $q$ -calculus operators in examinations of specific classes of functions which are analytic in  $\mathcal{U}$ . For further details on  $q$ -calculus one can refer to [13], [14], [15], and, [16], as well as the additional references mentioned therein.

Alongside the advancement of the theory and application of  $q$ -calculus, the theory of  $q$ -calculus dependent on two parameters  $(r, q)$ -integers has also been introduced and received more consideration during the last few decades. In 1991, [17], showed the  $(r, q)$ -calculus. Next, [18], investigated the fundamental theorem of  $(r, q)$ -calculus and a few  $(r, q)$ -Taylor formulas. As of late, [19], studied the  $(r, q)$ -derivative and  $(r, q)$ -integral on finite intervals. Besides, they concentrated on certain properties of  $(r, q)$ -calculus and  $(r, q)$ -associated of some important integral inequalities. The  $(r, q)$ -derivative have been considered and quickly created during this period by many creators.

Classes defined by derivative operators often arise in complex analysis or functional analysis and are crucial in characterizing specific sets of functions that satisfy certain properties related to their derivatives. These classes help classify functions based on their behavior under differentiation or specialized derivative operators. Utilizing the above defined  $(r, q)$ -calculus, certain subclasses in the class  $\hat{\mathcal{A}}(p)$  have as of now been explored in geometric function theory. Some classes defined by  $(r, q)$ -calculus operators like  $q$ -starlike and  $q$ -convex functions, these classes are defined using  $q$ -calculus operators. For instance,  $q$ -starlike functions are those for which the  $q$ -derivative has positive real part in the unit disk under the  $q$ -calculus framework. Similarly,  $q$ -convex functions satisfy certain conditions related to their  $q$ -derivatives in the unit disk. Also  $(r, q)$ -starlike and  $(r, q)$ -convex functions, these classes are extensions of the  $q$ -starlike and  $q$ -convex classes, incorporating additional parameters,  $r$  and  $q$  in the calculus operators.  $(r, q)$ -starlike and  $(r, q)$ -convex functions exhibit specific properties related to their derivatives under the  $(r, q)$ -calculus framework. [20], were the first who utilized the  $q$ -derivative operator  $\mathcal{D}_q$  to concentrate on the  $q$ -calculus comparable of the class

$S^*$  of starlike functions in  $\mathcal{U}$ .

Now, let  $\mathcal{M}(A, B, C)$  be the subclass of  $\hat{\mathbb{A}}(1)$  consisting of functions  $f \in \hat{\mathbb{A}}(1)$  which satisfy the inequality

$$\left| \frac{f'(z) - 1}{Af'(z) + (1 - B)} \right| < C,$$

where  $0 \leq A \leq 1$ ,  $0 \leq B < 1$  and  $0 < C \leq 1$  for all  $z \in \mathcal{U}$ . This class of functions was studied by, [21].

In, [21], defined the class  $\mathcal{M}^T(A, B, C)$  by  $\mathcal{M}^T(A, B, C) = \mathcal{M}(A, B, C) \cap \mathcal{T}$ .

Further we present some general subclass of analytic and multivalent functions related to  $(r, q)$ -derivative operator as follows.

**Definition 1.1** For  $0 \leq \alpha \leq 1$ ,  $0 \leq \beta < 1$ ,  $0 \leq \gamma < 1$ ,  $k \geq 0$ ,  $0 < q < r \leq 1$  and  $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ , we let  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  consist of functions  $f \in \mathcal{T}(P)$  satisfying the condition

$$\begin{aligned} & \operatorname{Re} \left( \frac{(\mathcal{D}_{r,q}f(z))' - 1}{\alpha(\mathcal{D}_{r,q}f(z))' + (1 - \gamma)} \right) \\ & > k \left| \frac{(\mathcal{D}_{r,q}f(z))' - 1}{\alpha(\mathcal{D}_{r,q}f(z))' + (1 - \gamma)} - 1 \right| + \beta. \end{aligned} \quad (3)$$

Certainly, the study outlined here appears to be focused on a specific class of mathematical functions denoted as  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ . The initial finding, the coefficient estimate or determine the coefficients of functions within the specified class, for functions  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ . These coefficients likely hold crucial information about the behavior and properties of these functions. Growth and distortion theorem are included, by including growth and distortion theorems, the study aims to investigate how these functions behave under certain transformations or conditions. Delving into their growth and distortion characteristics offers valuable insights into their behavior and potential practical uses. Furthermore, we obtain the extreme points, the study seeks to identify extreme points of these functions. These extreme points often hold significance in understanding the behavior and nature of the functions. After that, the radius of starlikeness and convexity, for the function in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  are determined. Understanding the boundaries within these functions possess such geometric properties is crucial in their characterization.

As a matter of first importance, let us take at

the coefficient inequalities. And the technique which studied in, [22], and also in [23].

## 2 Coefficient Inequalities

In this section we present a fundamental and sufficient condition for the function  $f$  in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ . Our main first result as follows:

**Theorem 2.1** Let  $0 \leq \alpha \leq 1$ ,  $0 \leq \beta < 1$ ,  $0 \leq \gamma < 1$ ,  $k \geq 0$ ,  $0 < q < r \leq 1$  and  $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ . A function  $f$  given by (1) is in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  if and only if

$$\sum_{n=p+1}^{\infty} \mu_n a_n \leq \mu_p + 1, \quad (4)$$

where

$$\mu_n = \left( \frac{(k+1)(1-\alpha) + \alpha(1-\beta)}{(k+\beta)(1-\gamma) + k + 1} \right) (n-1)[n]_{r,q}. \quad (5)$$

**Proof.** We have  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  if and only if the condition (3) is satisfied.

Let

$$w = \frac{(\mathcal{D}_{r,q}f(z))' - 1}{\alpha(\mathcal{D}_{r,q}f(z))' + 1 - \gamma},$$

upon the fact that,

$$\operatorname{Re}(w) \geq k|w - 1| + \beta \quad \text{if and only if} \\ (k+1)|w - 1| \leq 1 - \beta.$$

Now

$$\begin{aligned} & (k+1)|w - 1| \\ & = (k+1) \left| \left\{ \left( \sum_{n=p+1}^{\infty} (\alpha-1)[n]_{r,q}(n-1)a_n z^{n-2} \right) \right. \right. \\ & \quad - (\alpha-1)[p]_{r,q}(p-1)z^{p-2} - (2-\gamma) \left. \right\} \\ & \quad / \left\{ \alpha[p]_{r,q}(p-1)z^{p-2} \right. \\ & \quad \left. - \alpha \left( \sum_{n=p+1}^{\infty} [n]_{r,q}(n-1)a_n z^{n-2} \right) - (\gamma-1) \right\} \right| \\ & \leq 1 - \beta. \end{aligned} \quad (6)$$

The above inequality reduces to

$$\begin{aligned} & (k+1) \left( \left| \sum_{n=p+1}^{\infty} (\alpha-1)[n]_{r,q}(n-1)a_n z^{n-2} \right| \right. \\ & \quad - \left| (\alpha-1)[p]_{r,q}(p-1)z^{p-2} \right| - |2-\gamma| \left. \right) \\ & \quad / \left( \left| \alpha[p]_{r,q}(p-1)z^{p-2} \right| \right. \\ & \quad \left. - \alpha \left| \sum_{n=p+1}^{\infty} [n]_{r,q}(n-1)a_n z^{n-2} \right| - |\gamma-1| \right) \\ & \leq 1 - \beta. \end{aligned} \quad (7)$$

After that,

$$\begin{aligned} & (k+1) \left( \sum_{n=p+1}^{\infty} (1-\alpha) [n]_{r,q} (n-1) a_n \right. \\ & - (1-\alpha) [p]_{r,q} (p-1) - (2-\gamma) \Big) \\ & / \left( \alpha [p]_{r,q} (p-1) \right. \\ & - \alpha \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n - (1-\gamma) \Big) \\ & \leq 1-\beta. \quad \text{where } |z| < 1. \end{aligned}$$

then, we have

$$\begin{aligned} & \sum_{n=p+1}^{\infty} ((k+1)(1-\alpha) + \alpha(1-\beta)) (n-1) [n]_{r,q} a_n \\ & \leq ((k+1)(1-\alpha) + \alpha(1-\beta)) (p-1) [p]_{r,q} \\ & - (1-\gamma)(1-\beta) + (k+1)(2-\gamma). \end{aligned}$$

Thus

$$\begin{aligned} & \sum_{n=p+1}^{\infty} ((k+1)(1-\alpha) + \alpha(1-\beta)) (n-1) [n]_{r,q} a_n \\ & \leq ((k+1)(1-\alpha) + \alpha(1-\beta)) (p-1) [p]_{r,q} \\ & + (1-\gamma)(k+\beta) + (k+1), \end{aligned}$$

divide by  $(1-\gamma)(k+\beta) + (k+1)$  for both side, which yield to (4).

Suppose that (4) holds and we have to show (6) holds. Here the inequality (4) is equivalent to (7). So it suffices to show that,

$$\begin{aligned} & \left| \left\{ \left( \sum_{n=p+1}^{\infty} (\alpha-1) [n]_{r,q} (n-1) a_n z^{n-2} \right) \right. \right. \\ & - (\alpha-1) [p]_{r,q} (p-1) z^{p-2} - (2-\gamma) \Big\} \\ & / \left\{ \alpha [p]_{r,q} (p-1) z^{p-2} \right. \\ & - \alpha \left( \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n z^{n-2} \right) - (\gamma-1) \Big\} \Big| \\ & \leq \left\{ \left( \sum_{n=p+1}^{\infty} (1-\alpha) [n]_{r,q} (n-1) a_n \right) \right. \\ & - (1-\alpha) [p]_{r,q} (p-1) - (2-\gamma) \Big\} \\ & / \left\{ \alpha [p]_{r,q} (p-1) \right. \\ & - \alpha \left( \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n \right) - (1-\gamma) \Big\}. \quad (8) \end{aligned}$$

Since,

$$\begin{aligned} & \left| \alpha [p]_{r,q} (p-1) z^{p-2} - \left( \alpha \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n z^{n-2} \right) \right. \\ & - (\gamma-1) \Big|, \\ & \geq \left| \alpha [p]_{r,q} (p-1) z^{p-2} \right| - \left| \alpha \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n z^{n-2} \right| \\ & - |\gamma-1|, \end{aligned}$$

we have

$$\begin{aligned} & \left| \alpha [p]_{r,q} (p-1) z^{p-2} - \left( \alpha \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n z^{n-2} \right) \right. \\ & - (\gamma-1) \Big|, \\ & \geq \alpha [p]_{r,q} (p-1) - \alpha \sum_{n=p+1}^{\infty} [n]_{r,q} (n-1) a_n - (1-\gamma), \end{aligned}$$

where  $|z| < 1$ , and hence, we obtain (8).

**Theorem 2.2** Let  $0 \leq \alpha \leq 1$ ,  $0 \leq \beta < 1$ ,  $0 \leq \gamma < 1$ ,  $k \geq 0$ ,  $0 < q < r \leq 1$  and  $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ . If the function  $f$  given by (1) be in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  then

$$a_n \leq \frac{\mu_p + 1}{\mu_n}, \quad n = p+1, p+2, p+3, \dots, \quad (9)$$

where  $\mu_n$  is given by (5).

Equality holds for the functions  $f$  given by,

$$f(z) = z^p - \frac{(\mu_p + 1) z^n}{\mu_n}. \quad (10)$$

**Proof.** Since  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  Theorem 2.1 holds.

Now

$$\sum_{n=p+1}^{\infty} \mu_n a_n \leq \mu_p + 1,$$

we have,

$$a_n \leq \frac{\mu_p + 1}{\mu_n}.$$

Clearly the function given by (10) satisfies (9) and therefore  $f$  given by (10) is in  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  for this function, the result is clearly sharp.

### 3 Growth and Distortion Theorems for the Subclass $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$

The growth and distortion theorems represent foundational principles within complex analysis, focusing on comprehending how analytic functions behave,

grow, and affect geometric properties particularly in the context of univalent functions. These theorems play a crucial role in understanding the mappings and transformations of complex-valued functions. Both the growth and distortion theorems are especially important when dealing with univalent functions, which are functions that are injective or one-to-one within a certain domain. These theorems provide insights into the behavior of such functions and are crucial in various areas of mathematics and its applications.

The growth and distortion theorem will be considered and the covering property for function in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  is given by the following theorems.

**Theorem 3.1** Let  $0 \leq \alpha \leq 1, 0 \leq \beta < 1, 0 \leq \gamma < 1, k \geq 0, 0 < q < r \leq 1$  and  $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ . If the function  $f$  given by (1) be in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  then for  $0 < |z| = l < 1$ , we have

$$l^p - \frac{\mu_p + 1}{\mu_{p+1}} l^{p+1} \leq |f(z)| \leq l^p + \frac{\mu_p + 1}{\mu_{p+1}} l^{p+1}. \quad (11)$$

Equality holds for the function,

$$f(z) = z^p - \frac{\mu_p + 1}{\mu_{p+1}} z^{p+1}, \quad (z = \pm l, \pm il),$$

where  $\mu_p$  and  $\mu_{p+1}$  can be found by (5).

**Proof.** We only prove the right hand side inequality in (11), since the other inequality can be justified using similar arguments.

Since  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  by Theorem 2.1 we have,

$$\sum_{n=p+1}^{\infty} \mu_n a_n \leq \mu_p + 1.$$

Now

$$\begin{aligned} \mu_{p+1} \sum_{n=p+1}^{\infty} a_n &= \sum_{n=p+1}^{\infty} \mu_{p+1} a_n, \\ &\leq \sum_{n=p+1}^{\infty} \mu_n a_n \\ &\leq \mu_p + 1. \end{aligned}$$

And therefore

$$\sum_{n=p+1}^{\infty} a_n \leq \frac{\mu_p + 1}{\mu_{p+1}}, \quad (12)$$

since

$$f(z) = z^p - \sum_{n=p+1}^{\infty} a_n z^n,$$

we have,

$$\begin{aligned} |f(z)| &= \left| z^p - \sum_{n=p+1}^{\infty} a_n z^n \right|, \\ &\leq |z|^p + |z|^{p+1} \sum_{n=p+1}^{\infty} a_n |z|^{n-(p+1)}, \\ &\leq l^p + l^{p+1} \sum_{n=p+1}^{\infty} a_n. \end{aligned}$$

By aid of inequality (12), yields the right hand side inequality of (11).

**Theorem 3.2** If the function  $f$  given by (1) is in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  for  $0 < |z| = l < 1$  then, we have

$$\begin{aligned} p l^{p-1} - \frac{(p+1)(\mu_p + 1)}{\mu_{p+1}} l^p &\leq |f'(z)| \\ &\leq p l^{p-1} + \frac{(p+1)(\mu_p + 1)}{\mu_{p+1}} l^p. \end{aligned} \quad (13)$$

Equality holds for the function  $f$  given by

$$f(z) = z^p - \frac{\mu_p + 1}{\mu_{p+1}} z^{p+1}, \quad (z = \pm l, \pm il),$$

where  $\mu_p$  and  $\mu_{p+1}$  can be found by (5).

**Proof.** Since  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  by Theorem 2.1 we have

$$\sum_{n=p+1}^{\infty} \mu_n a_n \leq \mu_p + 1.$$

Now,

$$\begin{aligned} \mu_{p+1} \sum_{n=p+1}^{\infty} n a_n &\leq (p+1) \sum_{n=p+1}^{\infty} \mu_n a_n \\ &\leq (p+1)(\mu_p + 1). \end{aligned}$$

Hence

$$\sum_{n=p+1}^{\infty} n a_n \leq \frac{(p+1)(\mu_p + 1)}{\mu_{p+1}}, \quad (14)$$

since

$$f'(z) = p z^{p-1} - \sum_{n=p+1}^{\infty} n a_n z^{n-1}.$$

Then, we have

$$p |z|^{p-1} - |z|^p \sum_{n=p+1}^{\infty} na_n |z|^{n-1-p} \leq |f'(z)| \leq p |z|^{p-1} + |z|^p \sum_{n=p+1}^{\infty} na_n |z|^{n-1-p},$$

where  $|z| < 1$ . By using the inequality (14), we get Theorem 3.2 and this completes the proof.

**Theorem 3.3** If the function  $f$  given by (1) is in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  then  $f$  is starlike of order  $\delta$ , where

$$\delta = 1 - \frac{(\mu_p + 1)p}{-(\mu_p + 1) + \mu_{p+1}}.$$

The result is sharp with

$$f(z) = z^p - \frac{\mu_p + 1}{\mu_{p+1}} z^{p+1},$$

where  $\mu_p$  and  $\mu_{p+1}$  can be found by (5).

**Proof.** It suffices to show that (4) implies

$$\sum_{n=p+1}^{\infty} a_n(n - \delta) \leq 1 - \delta. \quad (15)$$

That is,

$$\frac{n - \delta}{1 - \delta} \leq \frac{\mu_n}{\mu_p + 1}, \quad n \geq p + 1. \quad (16)$$

The above inequality is equivalent to

$$\delta \leq 1 - \frac{(n - 1)(\mu_p + 1)}{-(\mu_p + 1) + \mu_n} = \psi(n),$$

where  $n \geq p + 1$ .

And  $\psi(n) \geq \psi(p + 1)$ , (16) holds true for any  $0 \leq \alpha \leq 1$ ,  $0 \leq \beta < 1$ ,  $0 \leq \gamma < 1$ ,  $k \geq 0$ ,  $0 < q < r \leq 1$  and  $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ . This completes the proof of Theorem 3.3.

#### 4 Extreme Points of the Class

$$\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$$

In spite of its elegance, the significance of extreme point theory within complex function theory is relatively limited. Several papers authored by Brickman, Hallenbeck, Mac Gregor, and Wilken have specifically determined extreme points within traditional families of analytic functions. A comprehensive

overview of their findings can be found in [24]. Notably, the availability of extreme points within the set  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ , comprising functions  $f$  that are analytic on the unit disc, possess a positive real part, and are normalized by  $f(0) = 1$ , holds fundamental importance. The theorem detailing the extreme points of the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  is as follows.

**Theorem 4.1** Let  $f_p(z) = z^p$ , and

$$f_n(z) = z^p - \frac{\mu_p + 1}{\mu_n} z^n, \quad n = p + 1, p + 2, p + 3, \dots,$$

where  $\mu_n$  is given by (5).

Then  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  if and only if it can be represented in the form

$$f(z) = \sum_{n=p}^{\infty} y_n f_n(z) \quad (17)$$

where  $y_n \geq 0$  and  $\sum_{n=p}^{\infty} y_n = 1$ .

**Proof.** Suppose  $f$  can be expressed as in (17). Our goal is to show that  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ .

By (17) we have

$$\begin{aligned} f(z) &= \sum_{n=p}^{\infty} y_n f_n(z) \\ &= y_p f_p(z) + \sum_{n=p+1}^{\infty} y_n f_n(z), \\ &= y_p z^p + \sum_{n=p+1}^{\infty} y_n \left( z^p - \frac{\mu_p + 1}{\mu_n} z^n \right), \\ &= y_p z^p + \sum_{n=p+1}^{\infty} y_n z^p - \sum_{n=p+1}^{\infty} y_n \frac{\mu_p + 1}{\mu_n} z^n, \\ &= z^p - \sum_{n=p+1}^{\infty} \frac{(\mu_p + 1)y_n}{\mu_n} z^n. \end{aligned}$$

After that,  $f(z) = z^p - \sum_{n=p+1}^{\infty} a_n z^n$  we see  $a_n = \frac{(\mu_p + 1)y_n}{\mu_n}$ ,  $n \geq p + 1$ .

Now, we have  $\sum_{n=p}^{\infty} y_n = y_p + \sum_{n=p+1}^{\infty} y_n = 1$  then

$$\sum_{n=p+1}^{\infty} y_n = 1 - y_p \leq 1.$$

Setting

$$\sum_{n=p+1}^{\infty} y_n \frac{\mu_p + 1}{\mu_n} \times \frac{\mu_n}{\mu_p + 1}$$

$$= \sum_{n=p+1}^{\infty} y_n = 1 - y_p \leq 1.$$

It follows from Theorem 2.1 that the function  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ .

Conversely, it suffices to show that

$$a_n = \frac{\mu_p + 1}{\mu_n} y_n.$$

Now we have  $f \in \Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  then by previous Theorem 2.2.

$$a_n \leq \frac{\mu_p + 1}{\mu_n}, \quad n \geq p + 1.$$

That is,

$$\frac{\mu_n a_n}{\mu_p + 1} \leq 1,$$

but  $y_n \leq 1$ .

Setting,

$$y_n = \frac{\mu_n a_n}{\mu_p + 1}, \quad n \geq p + 1.$$

Thus yields to the desired result and completes the theorem.

**Corollary 4.2** The extreme point of the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  are the function

$$f_p(z) = z^p,$$

and

$$f_n(z) = z^p - \frac{\mu_p + 1}{\mu_n} z^n, \quad n = p + 1, p + 2, p + 3, \dots,$$

where  $\mu_n$  is given by (5).

Finally, in this paper we consider the radius of starlikeness and convexity.

## 5 Radius of Starlikeness and Convexity

The radius of starlikeness and convexity for the function in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$  will also be considered.

**Theorem 5.1** If the function  $f$  given by (1) is in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ , then  $f$  is starlike of order  $\delta$  ( $0 \leq \delta < p$ ), in the disk  $|z| < R$  where

$$R = \inf \left[ \frac{\mu_n}{\mu_p + 1} \times \left( \frac{p - \delta}{n - \delta} \right) \right]^{\frac{1}{n-p}}, \quad (18)$$

where  $n = p + 1, p + 2, p + 3, \dots$ , and  $\mu_n$  is given by (5).

**Proof.** Here (18) implies

$$(\mu_p + 1)(n - \delta) |z|^{n-p} \leq \mu_n (p - \delta).$$

It suffices to show that

$$\left| \frac{z f'(z)}{f(z)} - p \right| \leq p - \delta,$$

for  $|z| < R$ , we have

$$\left| \frac{z f'(z)}{f(z)} - p \right| \leq \frac{\sum_{n=p+1}^{\infty} (n - p) a_n |z|^{n-p}}{1 - \sum_{n=p+1}^{\infty} a_n |z|^{n-p}}. \quad (19)$$

By aid of (9), we have

$$\left| \frac{z f'(z)}{f(z)} - p \right| \leq \frac{\sum_{n=p+1}^{\infty} \frac{(\mu_p + 1)(n - p) |z|^{n-p}}{\mu_n}}{1 - \sum_{n=p+1}^{\infty} \frac{(\mu_p + 1) |z|^{n-p}}{\mu_n}}.$$

The last expression is bounded above by  $p - \delta$  if.

$$\sum_{n=p+1}^{\infty} \frac{(\mu_p + 1)(n - p) |z|^{n-p}}{\mu_n}$$

$$\leq \left[ 1 - \sum_{n=p+1}^{\infty} \frac{(\mu_p + 1) |z|^{n-p}}{\mu_n} \right] (p - \delta),$$

and it follows that

$$|z|^{n-p} \leq \left[ \frac{\mu_n}{\mu_p + 1} \left( \frac{p - \delta}{n - \delta} \right) \right], \quad n \geq p + 1$$

which is equivalent to our condition (18) of the theorem.

**Theorem 5.2** If the function  $f$  given by (1) is in the class  $\Upsilon(\alpha, \beta, \gamma, k, r, q, p)$ , then  $f$  is convex of order  $\varepsilon$  ( $0 \leq \varepsilon < p$ ), in the disk  $|z| < w$  where

$$w = \inf \left[ \frac{\mu_n}{\mu_p + 1} \times \left( \frac{p(p - \varepsilon)}{n(n - \varepsilon)} \right) \right]^{\frac{1}{n-p}}$$

where  $n = p+1, p+2, p+3, \dots$ , and  $\mu_n$  is given by (5).

**Proof.** By using the same technique in the proof of Theorem 5.1, we can show that

$$\left| \frac{zf''(z)}{f'(z)} - (p-1) \right| \leq p - \varepsilon, \quad \text{for } |z| \leq w,$$

with the aid of (9). Thus we have the assertion of Theorem 5.2.

## 6 Conclusion

In this article, our primary focus lies in investigating a newfound subclass of multivalent analytic functions within the open unit disc, delineated by the application of Jackson's derivative operator. Our exploration begins with a thorough investigation by uncovering the essential criteria for functions falling into this category through the lens of Coefficients' Characterization. This approach flattens many of intriguing features, with notable highlights encompassing coefficient estimates, theorems on growth and distortion, identification of extreme points, determination of the starlikeness radius, and exploration of convexity within this unique subclass. It is clear from the analysis in this article that the use of the Jackson derivative operator is not limited only to this specific subclass, but also opens up ways to derive broader classes of multivalent analytical functions. During the analysis, the importance of extending this approach to studying the description of parameters for these broader classes was emphasized, thus enriching our understanding of the diverse mathematical landscapes governed by the interaction of functions and their distinctive properties. We are going to branch out into uncharted territories, pushing the boundaries of knowledge in the fascinating realm of multivalent analytic functions.

### References:

- [1] H. Silverman, Univalent functions with negative coefficients, *Proceedings of the American mathematical society*, **51**. 1, (1975), 109-116.
- [2] M. Kunt, I. Iscan, N. Alp, and M. Z. Sarikaya,  $(p, q)$ -Hermite-Hadamard inequalities and  $(p, q)$ -estimates for midpoint type inequalities via convex and quasi-convex functions, *Revista De La Real Academia De Ciencias Exactas Fisicas Y Naturales Serie A-Matematicas*, **112**, (2018), 969-992.
- [3] J. Prabseang, K. Nonlaopon, and J. Tariboon,  $(p, q)$ -Hermite-Hadamard inequalities for double integral and  $(p, q)$ -differentiable convex function, *Axioms*, **8**. 2, (2019), 1-10.
- [4] P. N. Sadjang, On the  $(p, q)$ -gamma and the  $(p, q)$ -beta functions, *arXiv preprint arXiv:1506.07394*, 2015.
- [5] H. M. Srivastava, Some generalizations and basic (or  $q$ -) extensions of the Bernoulli, Euler and Genocchi polynomials, *Applied Mathematics and Information Sciences*, **5**(3), (2011), 390-444.
- [6] M. Tunc, and E. Göv,  $(p, q)$ -Integral inequalities, *RGMA research report collection*, **19**, (2016), 1-13.
- [7] H. Kalsoom, M. Amer, M. D. Junjua, S. Has-sain, and G. Shahzadi, Some  $(p, q)$ -estimates of Hermite- Hadamard-type inequalities for coordinated convex and quasi convex functions, *Mathematics*, **7**(8). (2019), 1-22.
- [8] M. I. Alharayzeh, On a subclass of  $k$ -uniformly analytic and multivalent functions defined by  $q$ -calculus operator, *Far East Journal of Mathematical Sciences*, 132(1), (2021), 1-20.
- [9] F. H. Jackson, On a  $q$ -definite integrals, *The Quarterly Journal of Pure and Applied Mathematics*, **41**(1910), 193-203.
- [10] F. H. Jackson,  $q$ -difference equations, *American journal of mathematics*, **32**.4. (1910), 305-314.
- [11] F. H. Jackson, Xi—On  $q$ -functions and a certain difference operator, *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, **46**.2, (1909), 253-281.
- [12] S. Kanas, and D. Raducanu, Some subclass of analytic functions related to conic domains, *Mathematica slovacca*, **64**, no. 5, (2014) , 1183-1196.
- [13] S. Araci, U. Duran, M. Acikgoz, and H. M. Srivastava, A certain  $(p, q)$ - derivative operator and associated divided differences, *Journal of Inequalities and Applications*, (2016), 1-8.
- [14] A. Aral, V. Gupta, and R. P. Agarwal, Applications of  $q$ -calculus in operator theory, *Springer, New York*, 2013.
- [15] M. Govindaraj, and S. Sivasubramanian, On a class of analytic function related to conic domains involving  $q$ -calculus, *Analysis Mathematica*, **43**. (3), (2017), 475-487.
- [16] Z. Karahuseyin, S. Altinkaya, and S. Yalcin, On  $H_3(1)$  Hankel determinant for univalent functions defined by using  $q$ -derivative operator, *Tu Jhoothi Main Makkaar*, **9** (2017), No. 1, 25-33.



- [17] R. Chakrabarti, and R. Jagannathan, A  $(p, q)$ -oscillator realization of two parameter quantum algebras, *Journal of Physics A: Mathematical and General*, **24**(13), (1991), L711-L718.
- [18] P. N. Sadjang, On the fundamental theorem of  $(p, q)$ -calculus and some  $(p, q)$ -Taylor formulas, *Results in Mathematics*, **73** (2018), 1-21.
- [19] M. Tunc, and E. Göv, Some integral inequalities via  $(p, q)$ -calculus on finite intervals, *Filomat*, **35**. 5, (2021), 1421-1430.
- [20] M. E. H. Ismail, E. Merkes, and D. Styer, A generalization of starlike functions, *Complex Variables, Theory and Application: An International Journal*, **14**.1-4, (1990), 77-84.
- [21] H. S. Kim, and S. K. Lee, Some classes of univalent functions, *Math. Japon*, **32**(5), (1987), 781-796.
- [22] E. Aqlan, J. M. Jahangiri, and S. R. Kulkarni, New classes of  $k$ -uniformly convex and star-like functions, *Tamkang Journal of Mathematics*, **35**(3), (2004), 261-266.
- [23] M. I. Alharayzeh, and H.S. Alzboon, On a subclass of  $k$ -uniformly analytic functions with negative coefficients and their properties, *Nonlinear Functional Analysis and Applications*, Vol. 28, No. 2 (2023), pp. 589-599.
- [24] D.J. Hallenbeck, and T.H. Mac Gregor, *Linear Problems and Convexity Techniques in Geometric Function Theory*, Pitman, (1984).

#### Authorship contribution

Mamoun Harayzeh Al-Abbadi: Find the results, Analysis, Methodology, Writing—original draft, Supervision, Writing—review editing. Maslina Darus: review and check the results, Data curation, Investigation and Visualization. Habis S. Al-zboon: Investigation, Writing—review and editing.

#### Acknowledgments

The authors would like to express deepest thanks to the reviewers for their insightful comments on their paper.

#### Funding

No funding was received for conducting this study.

#### Conflicts of Interest

The authors have no conflicts of interest to declare that are relevant to the content of this article.