

Some Consequences of Mathematical Inaccuracies in Mechanics

EVELINA PROZOROVA

Mathematical-Mechanical Department, St. Petersburg State University,
Av. 28, Peterhof, 198504
RUSSIA

Abstract -: The article is devoted to the study of some violations of the known laws of mathematics and classical mechanics in continuum mechanics and kinetics. The most common is open non-stationary systems. From the equations formulated earlier and some experiments, a connection traced between the gradients of physical quantities and the angular momentum (force). The use of Hamilton's formalism and the dependence of force only on the distance between particles limits the study. In the collision integral, for example, for a rarefied gas, the Lennard-Jones potential, which not related to the type under consideration, is often used. Hamilton's formalism traces the behavior of closed systems. The general form of boundary conditions and forces changes the theory proposed in the works of N.N. Bogolyubov. The results of the reformulation discussed. Even in the classical theory, after taking into account the moments, we come to the absence of symmetric stress tensor in Boltzmann theory. The symmetric tensor obtains after assumption of small influence from absence of symmetry at the condition of the forces balance. No symmetric tensor leads to the existence of two solutions. New examples of solving problems on hydromechanics, elasticity theory and kinetic theory are given. A correspondence between the terms of the Liouville equation with more general and traditional forces established for continues mechanics. Previously considered boundary layer problems, jet problems and the simplest problems of elasticity theory. The paper proposes a method for finding the second solution for no symmetric problems, if the solution of the symmetric problem we know. The mathematical inaccuracies of the theory of continuum mechanics and kinetics discussed.

Keywords: Angular Momentum, Non-Symmetrical Stress Tensor, Hamiltonian formalism.

Received: June 16, 2021. Revised: June 25, 2022. Accepted: July 23, 2022. Published: September 16, 2022.

1 Introduction

If we consider that continuum mechanics "is a part of mathematics that supplies and investigates logical models for describing the changes in positions and forms that are undergone by things that we observe everyday" [1], then we can analyse it's as part of mathematics. The pitfall of this representation is the choice of a mathematical model, which obtained from a physical model. The physical model obtains from experiments or from previous theories. The experiment is done on the basis of material objects. Mathematical models differ from physical models by the definition of a point [2]. A point has no size, a physical object is always characterized by a size, at least by a wavelength. The rotational movement of the point does not make sense. Computational mathematics always deals with dimensional quantities. Differential equations obtained from integral equations under the additional condition that the functions are smooth, i.e. the space of piecewise continuous functions becomes the space of smooth

functions. Accordingly, there are no mathematical lines in physics. Representation in the form of points, lines and other infinitesimal quantities change the physical content of the problem. In mechanics, conservation laws were obtained in their derivation from integral equations under the assumption of an insignificant role outside the integral term [3]. Further simplification connected with the rejection of the tangential velocity component on the sides of an arbitrary elementary volume, which is a consequence of the rejection outside the integral term. This happens when designing, for example, density. Thus, any rotations of the elementary volume not allowed. The action of the moment is therefore was excluded. Any interactions excluded, except interactions along the normal. Comparison of the results of analysis for derivatives calculated through the distribution function and through the sum of the delta functions shows that the flows through the boundary are not take into account in the calculations through distribution function. Change of the system of particles only in the volume [4],[6],[7],[8] is considered.

Therefore, the method of molecular particles is more adequate for a large number of particles than the solution of the Boltzmann equation. Consideration of stochastic equations based on the sum of two integrals and the concept of a physically infinitesimal scale leads to the diffusion effect. Hilbert hypothesis about the time that dependence only from the distribution function through macroparameters confirmed, but becomes more complex. When deriving the Boltzmann equation, all inter particles correlations neglected, which is unfair when taking into account the action of the angular momentum associated with the movement of particles. The difference between the model that building on the basis the distribution function and by the method molecular dynamics, shows the hypothesis of molecular chaos remains, but the "Markovian" nature of the process is partially lost. Flows pass through the surfaces, which leads to the appearance of a diffusion flow not only in the collision integral, but also outside the collision integral. Flows depend on the history of their formation. These flows are not dissipative. Let us dwell on the models used in more detail. The most common systems are open non-stationary systems. Accumulation of experimental facts led to the hypothesis of the importance of spatial gradients. From the record of the modified equations and some experiments, the connection between the gradients of physical quantities and the momentum under consideration are traced. Being an open system, the elementary volume exchanges components of physical quantities not only along the normal, but also along the tangential direction. Mathematical modeling of such objects causes difficulties, since their description is represented by a system of non-linear non-stationary equations. Previous works have shown that classical models do not include one of the most important laws - the law of conservation of angular momentum, if the moment does not act as a given force effect [4],[5],[6],[7],[8],[9]. When the action of the distributed moment is taken into account, the system of equations is supplemented by the moment conservation law. In the classical theory, since the symmetry of the stress tensor is postulated, the law is identically satisfied due to the vanishing of the sum of stresses under equilibrium conditions. The paper shows that the no symmetric tensor corresponds to two types of solutions. As a rule, the analysis of the mathematical properties of such systems is performed without analyzing the role of the initial and boundary conditions. So, for example, it is difficult to choose a problem that satisfies the group properties of the equations of gas dynamics,

elasticity theory, kinetic equations, etc. The self-similar solutions obtained from the stretch group are, in fact, a consequence of singular initial conditions. In practice, the required conditions can rarely be met. Thus, the number of groups used in practice is limited not only by the properties of the equations, but also by the initial and boundary conditions. The same is true for closed and open systems. The performed mathematical analysis of the equations of continuum mechanics with the no symmetric stress tensor showed that in the plane case for four unknowns in the classical formulation we have three equations: two equations from the stress equilibrium condition and one equation - the moment equilibrium condition. Thus, we need to close the problem with an additional condition. In the classical version, such a condition is the condition of symmetry of the stress tensor. From definition of pressure, both from the classical Boltzmann equation and modified one, does not follow that the hydrostatic pressure is one third of the sum of the pressures on the coordinate areas. Using Pascal's law for equilibrium, we choose a pressure equal to one third of the pressures on the coordinate sites. However, the theory remains the same when determining different pressures at each of the sites, i.e. p_x, p_y, p_z . The use of one pressure is possible under equilibrium conditions (Pascal's law), but for non-equilibrium conditions the fact is not obvious. Neglecting the integral term when taking integrals by parts (the Ostrogradskii-Gauss theorem) is possible only for slow laminar flows. The different pressure is indicated by the analysis of the determination of pressures when comparing the results of its determination through potential velocities by substitution into the Euler equations. Writing separately the law of equilibrium for forces and separately for moments of forces without taking into account the mutual influence, although the moment creates an additional force, we come to the conclusion about the symmetry of the stress tensor. If we consider equal pressures in different directions, we lose the moment of force but the pressure gradient is a force. Analyzing the results of solving the Euler equations [6][7] and calculating potential flows, we obtain a vortex sheet, which indicates the existence of a moment. The numerical results were processed by the authors without averaging the values over the sides of the elementary volume. An analysis of the correspondence between the solutions for the potential flow and for the Euler equations indicates their no coincidence. The solution of two-dimensional problems means a special distribution of forces in the third dimension. In stochastic

processes of open systems, the motion of fast molecules is accompanied by a change in the position of the center of inertia, which is accompanied by the appearance of a moment. The appearance of torque leads to a change in the direction of velocities and the formation of local structures. It can be assumed that the moment will make a significant contribution to the equation of state of liquids and gases at high pressures (virial equation), since even for equilibrium distribution functions there are fast and slow molecules. It is possible to obtain a solution to the problem with an asymmetric tensor by applying an iterative procedure.

To do this, you must first solve the problem with a symmetric tensor, leave one solution obtained, and then use the second equation and solve it as a differential one. This would be one of the possible solutions. In three-dimensional problems, three stress equilibrium equations and three equations for moments are solved. In the classical version the symmetry of stress tensor condition again closes the problem. The algorithm for solving a three-dimensional problem is the same as for a two-dimensional one. The class of systems considered in statistical mechanics textbooks includes systems described by Hamiltonian dynamics. In the modern version, this class includes all systems described by the laws of classical or quantum mechanics. Since in our theory an important role is assigned to the law of conservation of the moment of force (momentum), it is necessary to study how the properties and structure of the system change when using the law of conservation of moment, since it was not included in statistical theories. We have already noted that a feature of classical theories is the identification of the properties of systems without studying the influence of boundary and initial conditions on their state. The basic equation in kinetic theory is the Liouville equation for a closed system. As already noted, the mathematical apparatus for deriving the equation is Hamilton's formalism. Questions arise about changing the character of equations for open systems. The definition of the distribution function of an open system for a non-stationary problem will depend on the initial and boundary conditions. For example, in the theory of N. N. Bogolyubov, boundary conditions are set with the requirement that the derivatives of the distribution function vanish at infinity. A special type of force action $\Phi(r_i - r_j)$ is used [2][10]. On this basis, a number of terms are assumed to be equal to zero. When considering flying aircraft, missiles, the boundaries of the perturbation region are limited and the behavior of

the generalized coordinates and impulses is different. Perturbations can also exist at infinity. An important component is the circulation component of the velocity, which ceases to be potential and additional dissipation occurs. The resulting stresses create a global circulation motion around the apparatus and lead to beats during movement, especially in the case of no symmetric design. To illustrate the action of the moment, model problems of fluid mechanics and the theory of elasticity are considered. For the problem of the first type, a self-similar problem was chosen, which was solved by A.G. Petrov, in the classical formulation [11]: Exact solution of the Navier-Stokes equations in a liquid layer between plates moving in parallel. As a second example, we consider the problem of a concentrated force F on a half-space at an arbitrary point with coordinates x, z . Correspondence between individual force terms of a general form and terms in the Liouville and Boltzmann equations and the equations of aeromechanics is clarified. Thus, the article demonstrates the contribution of the moment to the processes of aeromechanics and the theory of elasticity. The local interaction of molecules leads to the collective behavior of particles. From the point of view of statistics, the large size of the system turns out to be useful, since then it is possible to consider the average values of physical quantities. The trajectory of molecules with a small number of particles turns out to be confusing. However, the statistical description does not include the action of some factors, for example, the action of flows at the border. Correspondence between individual force terms of a general form and terms in the Liouville and Boltzmann equations and the equations of aeromechanics is clarified. Thus, the article demonstrates the contribution of the moment to the processes of aeromechanics, the theory of elasticity and kinetics. In fact, questions of the influence of non-conservative forces and their contribution to macroequations are discussed. The results of the work consist in illustrating the influence of the moment on the example of solving the problems of fluid and solid mechanics, developing a general methodology for determining a non-symmetric operator if a solution with a symmetric operator is found, constructing a more general kinetic theory based on the Liouville equation with a complicated type of acting forces; analysis of mathematical models of continuum mechanics and kinetic theory in order to refine the accepted classical models.

2. Examples of Solution Equations with Influence Angular Momentum

a. Fluid between parallel moving plates. In previous works, the general equations of continuum mechanics were obtained taking into account совместное the influence of the angular momentum and force. There are few analytical solutions of viscous fluid problems, even two-dimensional ones. For one-dimensional problems, the moment does not work. Therefore, we restrict ourselves to the simplest two-dimensional ones. Consider, as an example, the unsteady motion of a fluid between two moving plates. A.G. Petrov showed the self-similarity of the problem in the classical formulation and obtained an analytical solution [11]. Let us try to determine the influence of the no symmetry of the stress tensor. So far with the old density equation.

$$u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = -\frac{\partial p}{\partial x} + \nu \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right),$$

$$u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} = -\frac{\partial p}{\partial y} + \nu \left(\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} \right) + \nu \frac{\partial}{\partial y} \left(y \frac{\partial^2 v}{\partial y^2} \right), \quad (1)$$

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial x} = 0,$$

$$x \left(\frac{\partial \sigma_{xz}}{\partial x} + \frac{\partial \sigma_{zz}}{\partial z} \right) - z \left(\frac{\partial \sigma_{xx}}{\partial x} + \frac{\partial \sigma_{zx}}{\partial z} \right) + \sigma_{zx} - \sigma_{xz} = 0.$$

With boundary conditions

$$u(t, x, 0) = u(t, x, h) = 0, \quad v(t, x, 0) = 0, \quad v(t, x, h) = \dot{h}.$$

ν - kinematic viscosity, the density is considered equal to one.

u, v - speeds in x, y coordinates. p - pressure. The equations differ from those of A.G. Petrov by the last term, which describes the action of the moment. We are looking for a new solution, leaving a functional depend About the kinetic description.

$$v(t, x, h) = \dot{h}V(\eta), \quad u = \frac{\dot{h}}{h} xU(\eta), \quad p = \dot{h}^2 \left[b \frac{x^2}{2h^2} + P(t, \eta) \right] + p_0, \quad \eta = \frac{y}{h}. \quad (2)$$

After substitution, we get

$$U + V' = 0, \quad -aU - \eta U' + U^2 + V U' + b - \frac{1}{Re} U'' + \frac{1}{Re} (\eta U'')' = 0. \quad (3)$$

$$U(0) = V(0) = 0, \quad U(1) = 0, \quad V(1) = 1, \quad a(t) = 1 - \frac{\dot{h}h}{\dot{h}^2}, \quad Re(t) = \frac{h\dot{h}}{\nu}. \quad (4)$$

In the case of plate motion according to the law $h = k\sqrt{|t - t_0|}$ we get

$$P = \eta V - \frac{V^2}{2} + \frac{1}{Re} V'(\eta). \quad (5)$$

$$u = \frac{\dot{h}}{h} x \left[\cos \left(2n\pi \frac{y}{h} \right) - 1 \right], \quad (6)$$

$$v = \dot{h} \left[\eta - \frac{1}{2\pi n} \sin(2\pi n \eta) \right] \quad (7)$$

Let's leave one old solution and define voltage difference

$$x \left(\frac{\partial \sigma_{xy}}{\partial x} + \frac{\partial \sigma_{yy}}{\partial y} \right) - y \left(\frac{\partial \sigma_{xx}}{\partial x} + \frac{\partial \sigma_{yx}}{\partial y} \right) + \sigma_{yx} - \sigma_{xy} = 0$$

$$u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} = -\frac{\partial p}{\partial x} + \nu \left(\frac{\partial \sigma_{xx}}{\partial x} + \frac{\partial (\sigma_{yx} + \Delta_{xy})}{\partial x} \right). \quad (8)$$

$$u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} = -\frac{\partial p}{\partial y} + \nu \left(\frac{\partial (\sigma_{yx} + \Delta_{xy})}{\partial x} + \frac{\partial \sigma_{yy}}{\partial x} \right),$$

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial x} = 0$$

Then, leaving the old solution along the x axis, let's look at the value of the addition to the no symmetric tensor.

$$x \left(\frac{\partial (\sigma_{yx} + \Delta_{xy})}{\partial x} + \frac{\partial \sigma_{yy}}{\partial y} \right) - \Delta_{xy} = 0.$$

$$x \frac{\partial \sigma_{yx}}{\partial x} + x \frac{\partial \Delta_{xy}}{\partial x} - \Delta_{xy} = 0.$$

$$\frac{\partial \Delta_{xy}}{\partial x} - \frac{1}{x} \Delta_{xy} + \frac{\partial \sigma_{yx}}{\partial x} = 0.$$

$$\frac{\partial \sigma_{yx}}{\partial x} = -\frac{\dot{h}}{h} \sin \left(2n\pi \frac{y}{h} \right) \quad (9)$$

$$\Delta_{xy} = e^{-\int \frac{1}{x} dx} \left[-\int \frac{\partial \sigma_{yx}}{\partial x} e^{\int \frac{1}{x} dx} dx + C \right], \quad \Delta_{xy} = \frac{1}{x} \int \frac{\dot{h}}{h} \sin \left(2n\pi \frac{y}{h} \right) \cdot x dx$$

Here Δ_{xy} is the difference between $\sigma_{zx} - \sigma_{xz}$. The solution of the new equation differs slightly in the coefficients at η . New values are determined using iterations. Most likely, the moment along the x axis ($\nu \frac{\partial}{\partial y} \left(y \frac{\partial^2 u}{\partial x^2} \right)$) plays the greatest role. Calculations are done in the same way, but more iterations are required.

b. Examples of the influence of the moment in the theory of elasticity. The problems are taken from the book [12]. Consider the problem when a concentrated force is given on a half-plane. Let us find the stress distribution with and without taking

into account the influence of the moment for problem (Fig. 1).

General view of equations

$$\frac{\partial \sigma_x}{\partial x} + \frac{\partial \sigma_{zx}}{\partial z} = 0, \quad \frac{\partial \sigma_z}{\partial z} + \frac{\partial \sigma_{xz}}{\partial x} = 0, \quad (10)$$

$$x \left(\frac{\partial \sigma_{xz}}{\partial x} + \frac{\partial \sigma_{zz}}{\partial z} \right) - z \left(\frac{\partial \sigma_{xx}}{\partial x} + \frac{\partial \sigma_{zx}}{\partial z} \right) + \sigma_{zx} - \sigma_{xz} = 0.$$




Fig. 1: The problem formulation of example.

Suggested that $\sigma_{zx} = -A \frac{x}{x^2 + z^2}$, option for case $\sigma_{xz} = \sigma_{zx}$ and $\sigma_z = -A \frac{z}{(x^2 + z^2)^2}$ was considered in [12]. Then, we solve the problem when condition of the stress tensor is satisfied $\sigma_{xz} \neq \sigma_{zx}$

$$\begin{aligned} \frac{\partial \sigma_{zx}}{\partial z} &= -A \frac{2xz}{(x^2 + z^2)^2}, \\ \sigma_x &= -2Az \int_0^x \frac{x dx}{(x^2 + z^2)^2} - Az \int_0^y \frac{dy}{(y + z^2)^2} \\ &= Az \frac{1}{(z^2 + y)} + f(z), \\ \sigma_x &= Az \frac{1}{(z^2 + x^2)} + f(z). \end{aligned} \quad (11)$$

$$\begin{aligned} \frac{\partial \sigma_{xz}}{\partial x} &= -A \left(\frac{1}{x^2 + z^2} - \frac{2x^2}{(x^2 + z^2)^2} \right) \\ &= -A \left(-\frac{z^2 - x^2}{(x^2 + z^2)^2} \right), \\ \sigma_z &= -A \int_0^z \left(\frac{1}{x^2 + z^2} - \frac{2x^2}{(x^2 + z^2)^2} \right) dz = \\ &= -A \frac{1}{x} \arctg z - A \left(\frac{1}{x} \frac{z}{1+(z^2)} + \frac{1}{x} \arctg z \right) = \\ &= -A \frac{1}{x} \frac{z}{1+(z^2)} + f(x). \end{aligned} \quad (12)$$

Let σ_x and σ_{zx} remain the same. Then from the equation for the moments it is possible to determine the difference a new value σ_{xz}

$$\frac{\partial \sigma_{xz}}{\partial x} - \frac{1}{x} \sigma_{xz} + \frac{\partial \sigma_{zz}}{\partial z} = 0.$$

$$x \left(\frac{\partial \sigma_{xz}}{\partial x} + \frac{\partial \sigma_{zz}}{\partial z} \right) + \sigma_{zx} - \sigma_{xz} = 0. \quad (13)$$

Then a new value σ_z ($\sigma_z = \sigma_{zz}$) determined. An example is given in order to demonstrate the many possible solutions in the two-dimensional case. Consequently, the solution of the two-dimensional problem under the condition of symmetry of the stress tensor is only one of the options to solve the problem. Algorithm for finding stresses with no symmetric tensor easily programmed using previously obtained results.

3 About Kinetic Theory

The macroequations of continuum mechanics, the special cases of which we have considered, can theoretically be obtained from the Boltzmann kinetic equation, which, in turn, can be derived from the Liouville equation. In practice, continuum equations are obtained experimentally. Since in the experiment we work with material objects, and not with points, the corresponding experimental data can be represented in integral form. After taking moments and averaging, we arrive at the equations of continuum mechanics. However, in the integral formulation, the individual components of the forces and their contribution to the individual terms of the equations are not single out. Mathematical translation equations from integral form to different is means the transition from a piecewise continuous space to a continuous one. In addition, the use of the Hamiltonian formalism and the dependence of the force only on the distance between particles limits the study; in the collision integral, for example, for a rarefied gas, the Lennard-Jones potential is often used, which does not belong to the type under consideration. The foregoing makes us to the study of the influence of forces of another type on equations of mechanics. Let us consider new members for force:

$$\begin{aligned} \mathbf{M} &= \sum_i m_i [\mathbf{x}_i, \dot{\mathbf{x}}_i]. \text{ In the most frequently studied case } U = U(x_i - x_j) \\ \frac{d\mathbf{M}}{dt} &= -\sum_{ij} \frac{dU_{ij}}{dr_{ij}} \frac{[\mathbf{x}_i, \mathbf{x}_j]}{r_{ij}}, \quad L = T - U, \\ \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_k} - \frac{\partial L}{\partial q_k} &= 0, \end{aligned} \quad (14)$$

T - kinetic energy, U - potential.

Usually the Hamiltonian contains only potential forces. Actually generalized forces represent the sum of forces [13]

$$Q = -B - Cq, \quad \dot{q}_i = 0, \quad B = D + \Gamma, \quad D^T = D, \quad \Gamma^T = -\Gamma \quad (15)$$

Forces $-Dq$ dissipative, Gq –gyroscopic,

$$-\Gamma\dot{q} = \frac{d}{dt} \frac{\partial V}{\partial \dot{q}} - \frac{\partial V}{\partial q}, \quad V(q, \dot{q}) = -\frac{1}{2} q^T \Gamma q. \quad (16)$$

The matrix of positional forces C also decomposes into the sum of symmetric and skew-symmetric parts..

$$C = K + N, \quad K^T = K, \quad N^T = -N \quad (17)$$

Conservative forces have potential

$$-Kq = \frac{\partial U}{\partial q}, \quad U = -\frac{1}{2} q^T K q. \quad (18)$$

Consequently, $L = T - Q_r$, where the latter is the work of the generalized forces.

Formally, one can define, as in classical mechanics

$$\dot{q}_i = \frac{\partial H(q, \dot{q})}{\partial \dot{q}_i}, \quad \ddot{q}_i = -\frac{\partial H(q, \dot{q})}{\partial q_i},$$

However, they often work with generalized velocities

$$(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_n) = (q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_l, p_{l+1}, \dots, p_n), \quad p_k = \frac{\partial T}{\partial \dot{q}_k} = \dot{q}_k \quad (19)$$

$$H = \sum_{1 \leq i \leq n} \left(\frac{p_i^2}{2m} + U(r_i, t) \right) + \frac{1}{2} \sum_{1 \leq i, j \leq n, i \neq j} \Phi(|r_i - r_j|). \quad (20)$$

Thus, in statistical mechanics, a particular case of forces with a Coulomb potential is considered. An interesting feature of modern kinetic theories is the use of other potentials in the integral. For example, Lennard-Jones potential. In this case, the acting force is the derivative of a complex function. The presence of other molecules j is partially taken into account. Another assumption is the potential nature of the action of circulation forces. This discards the dissipation of the finely vortex motion in the theory of turbulence. In addition, the action of the moment of forces arising from the movement of molecules is not taken into account. Thus, modern representations require a revision of the equations and the inclusion in the theory of initial and boundary conditions.

4 The Effect of Dispersion in the Boltzmann Equation

Consider the equations for the s -particle modified distribution function F_s [8]

$$\begin{aligned} \frac{1}{V^s} \frac{\partial F_s}{\partial t} = & -\frac{1}{V^s} \sum_{i=1}^s \frac{p_i}{m} \left(\frac{\partial F_s}{\partial r_i} + \frac{\partial}{\partial r_i} r_j \frac{\partial F_s}{\partial r_j} \right) \\ & + \frac{1}{2V^s} \sum_{\substack{i,j=1 \\ (j \neq i)}}^s \frac{\partial Q}{\partial r_i} \frac{\partial F_s}{\partial \xi_i} + \\ & + \frac{1}{V^{s+1}} \sum_{i=1}^s \int \sum_{j=s+1}^N \frac{\partial Q}{\partial r_i} \\ & \frac{\partial F_{s+1}(t, r_1, \dots, r_s, r_j, \xi_1, \dots, \xi_s, \xi_j)}{\partial p_i} dr_j d\xi_j + \quad (21) \\ & + \frac{1}{V^{s+1}} \sum_{s+1}^N \int \frac{\xi_j}{m} \frac{\partial F_{s+1}(t, r_1, \dots, r_s, r_j, \xi, \dots, \xi_s, \xi_j)}{\partial r_j} dr_j d\xi_j + \\ & + \frac{1}{V^{s+1}} \sum_{i=1}^s \sum_{j=s+1}^N \int \frac{\partial Q}{\partial r_j} dr_j d\xi_j \cdot \\ & \frac{\partial F_{s+1}(t, r_1, \dots, r_s, r_j, \xi, \dots, \xi_s, \xi_j)}{\partial p_j} \\ & + \frac{1}{2V^{s+2}} \sum_{\substack{i,j=s+1 \\ (j \neq i)}}^s \int \frac{\partial Q}{\partial r_i} \cdot \\ & \frac{\partial F_{s+2}(t, r_1, \dots, r_s, r_j, \xi_1, \dots, \xi_s, \xi_j)}{\partial p_j} \\ & \cdot dr_i dr_j d\xi_i d\xi_j \end{aligned}$$

It is usually assumed that when $|r_j| \rightarrow \infty \frac{\partial}{\partial r_j} F_n \rightarrow 0, |\xi_j| \rightarrow \infty \frac{\partial}{\partial \xi_j} F_n \rightarrow 0$

This is possible only for the equilibrium state at infinity. Otherwise, perturbations of the distribution function will create flows and form a circulation motion around the apparatus, giving an additional force. Such a situation is possible when the apparatus enters the environment created by another flying body. If we trace the correspondence of the terms of the Liouville equation to the terms Boltzmann equations, and then macroequations, it turns out that the Euler equation, taking into account the moment for pressure, corresponds to all the potential forces included in Q .

a. Exact solution for kinetic theory. The barometric Boltzmann formula. Gas is at stationary condition in field of force which have potential φ (this is analogy tasks from [14]), no taking into account the moment (second term)

$$\xi_i \frac{\partial f}{\partial x_i} + \xi_i \frac{\partial}{\partial x_i} x_i \frac{\partial f}{\partial x_i} - \frac{1}{m} \frac{\partial \varphi}{\partial x_i} \frac{\partial f}{\partial \xi_i} = J(f, f). \quad (22)$$

As before solution we shall be look for as

$f = A(x)e^{-B(x)\xi^2}$. We receive the old result $B = Const$. For $A(x)$ we have equation

$$\frac{dA}{dx} + \frac{d}{dx_i} x_i \frac{dA}{dx_i} + 2 \frac{AB}{m} \frac{\partial \varphi}{\partial x_i} = 0.$$

We have old result that is common for one-dimension tasks:

$$f = n_0 \left(\frac{m}{2\pi kT}\right)^{3/2} e^{-\frac{\varphi}{kT}} e^{-\frac{m}{2kT}\xi^2}.$$

General local-Maxwell distribution без учета момента is $f =$

$$n \left(\frac{m}{2\pi kT}\right)^{3/2} \exp\left\{-\frac{m}{2kT}c^2\right\}, \quad c = \xi - u.$$

Modification the Boltzmann equation is

$$\xi_i \frac{\partial f}{\partial x_i} + \xi_i \frac{\partial}{\partial x_i} x_i \frac{\partial f}{\partial x_i} - g_i \frac{\partial f}{\partial \xi_i} = J(f, f). \quad g = X/m \quad \text{-acceleration of molecules.}$$

Let local-Maxwell solution of the equation will be considered as in old algorithm

$$lnf = \gamma_0 + \gamma_i \xi_i + \gamma_4 \xi^2. \quad (23)$$

Then we obtain old equation and the changing one without viscosity and thermal conductivity.

$$\begin{aligned} \frac{\partial \gamma_0}{\partial t} + g_i \gamma_i &= 0, \\ \frac{\partial \gamma_i}{\partial t} + 2g_i \gamma_4 + \frac{\partial \gamma_0}{\partial x_i} + \frac{\partial \gamma_0}{\partial x_i} + \frac{1}{2} x_i \frac{\partial \gamma_0^2}{\partial x_i^2} + \\ \frac{\partial}{\partial x_i} x_i \frac{\partial \gamma_0}{\partial x_i} &= 0, \end{aligned} \quad (24)$$

$$\begin{aligned} \frac{\partial \gamma_4}{\partial t} \delta_{ij} + \frac{1}{2} \left(\frac{\partial \gamma_i}{\partial x_j} + \frac{\partial \gamma_j}{\partial x_i}\right) + \frac{1}{2} \left(\frac{\partial \gamma_i}{\partial x_j} + \frac{\partial \gamma_j}{\partial x_i}\right) + \\ + \frac{1}{2} * \frac{1}{2} (x_i + x_j) \left(\frac{\partial \gamma_i}{\partial x_j} + \frac{\partial \gamma_j}{\partial x_i}\right) \frac{\partial \gamma_0}{\partial x_i} + \\ \frac{1}{2} (x_i + x_j) \frac{1}{2} \left(\frac{\partial}{\partial x_j} \left(\frac{\partial \gamma_i}{\partial x_j} + \frac{\partial \gamma_j}{\partial x_i}\right) + \right. \\ \left. \frac{\partial}{\partial x_i} \left(\frac{\partial \gamma_i}{\partial x_j} + \frac{\partial \gamma_j}{\partial x_i}\right)\right) = 0, \quad \text{As before} \end{aligned}$$

$$\frac{\partial \gamma_4}{\partial x_i} = 0, \quad T = const.$$

As the result we shall receive modified gas-dynamic equations.

5 Conclusion

The article investigates some violations of the known laws of mathematics and classical mechanics in continuum mechanics and kinetics. The focus is on open non-stationary systems. The consequences arising in mechanics are discussed, when discarding outside the integral term when applying Ostrogradskii-Gauss theorem. The use of the Hamiltonian formalism and the dependence of the force complex, even if it depends only on the distance between particles, limits classical studies. In the collision integral, for example, for a rarefied

gas, the Lennard-Jones potential, which does not relate to the Hamiltonian type under consideration, is often used. Hamilton's formalism traces the behavior of closed systems. The general form of boundary conditions and forces change the theory, proposed in the works of N.N. Bogolyubov. Even in the classical theory, after taking into account the moments, we come to the absence of symmetric stress tensor in Boltzmann theory. The symmetric tensor obtains after assumption of small effect of the absence of symmetry and from the condition of the balance of forces. No symmetric tensor leads to the existence of two solutions. New examples of solving problems of hydromechanics, elasticity theory and kinetic theory are given. A correspondence between the terms of the Liouville equation with more general and traditional forces is established. Previously considered boundary layer problems, jet problems and the simplest problems of elasticity theory. The paper proposes a method for finding the second solution. The mathematical inaccuracies of the theory of continuum mechanics and kinetics are discussed. To validate the proposed model we need additional experimental and theoretical research.

Reference:

- [1] T.O. Truesdell. *Initial course of rational continuum mechanics*. M.: Mir.1975
- [2]. Yu.L. Klimontovich. *Statistical theory of open systems*. M.: LENNARD, 2019. 1. 524
- [3] L.G. Loytsyansky. *Mechanics of liquid*
- [4]. Evelina Prozorova. The Effect of Angular Momentum and Ostrogradskii-Gauss Theorem in the Equations of Mechanics. Wseas Transaction on Fluid Mechanics DOI: 11.37394/232013.2020.15.2 12.
- [5] Evelina Prozorova. The influence of the no symmetric stress tensor on the flow separation. Wseas Transaction on Applied and Theoretical Mechanics DOI: 10.37394/232011.2020.15.9
- [6]. Evelina Prozorova Consequences of the Ostrogradskii-Gauss theorem for numerical simulation in aeromechanics. Elsevier Granthaalayah, June 2020. 8. 6.
- [7]. Evelina Prozorova. Influence of angular momentum and dimension on dynamics in mechanics. Wseas Transaction on Applied and Theoretical Mechanics. DOI: 10.37394/232011.2021.16.1
- [8]. Evelina Prozorova. Mechanism of Formation for Fluctuation Phenomena. Proceeding

conference CHAOC-20. .729-736
<http://www.cmsim.org/chaos2021.html>

- [9].Oleg Galaev, Evelina Prozorova. Dispersion Effects in the Falkner-Skan Problem and in the Kinetic Theory. *Journal of Applied Mathematics and Physics*, 2017, 5, 522-537.
- [10].R.Balescu. *Equilibrium and nonequilibrium statistic mechanics*. A Wiley-Intersciences Publication John Willey and Sons. New-yourk-London...1975.406
- [11] A.G. Petrov. On exact solutions of the Navier-Stokes equations in a liquid layer between parallel moving plates. *Proceedings of the XXII International Conference on Computational Mechanics and Modern Applied Software Systems (VMSPPS'2021)*, September 4–13, Alushta.. M.: MAI Publishing House, 2021. 57-59
- [12]. E. A. Bulanov. *Moment stresses in the mechanics of solid, bulk and liquid bodies*. M.: University book. 2012. 140.
- [13].V.F. Zhuravlev, A.G. Petrov, M.M. Shunderyuk. *Selected Problems of Hamiltonian Mechanics*. M.:LENAND, 2018. 304.
- [14].M.N. Kogan. *Dynamics of rarefied gas*. M.: Science, 1967. 440.

Contribution of Individual Authors to the Creation of a Scientific Article (Ghostwriting Policy)

The author(s) contributed in the present research, at all stages from the formulation of the problem to the final findings and solution.

Sources of Funding for Research Presented in a Scientific Article or Scientific Article Itself

No funding was received for conducting this study.

Conflict of Interest

The author(s) declare no potential conflicts of interest concerning the research, authorship, or publication of this article.

Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

https://creativecommons.org/licenses/by/4.0/deed.en_US